

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

cc

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁵ :C07D 239/95, 239/96, 401/04, 405/04,
409/04, A01N 43/54

A1

(11) International Publication Number:

WO 94/26722

(43) International Publication Date: 24 November 1994 (24.11.94)

(21) International Application Number: PCT/US94/04965

(22) International Filing Date: 10 May 1994 (10.05.94)

(30) Priority Data:

08/060,629	12 May 1993 (12.05.93)	US
08/144,904	28 October 1993 (28.10.93)	US

(60) Parent Applications or Grants

(63) Related by Continuation

US	08/144,904 (CIP)
Filed on	28 October 1993 (28.10.93)
US	08/060,629 (CIP)
Filed on	12 May 1993 (12.05.93)

(71) Applicant (for all designated States except US): E.I. DU PONT
DE NEMOURS AND COMPANY [US/US]; 1007 Market
Street, Wilmington, DE 19898 (US).


(72) Inventors; and

(75) Inventors/Applicants (for US only): BEREZNAK, James,
Francis [US/US]; 153 Nottingham Court, Aston, PA
19014 (US). CHANG, Zen-Yu [CN/US]; 23 Staten Drive,
Hockessin, DE 19707 (US). SELBY, Thomas, Paul
[US/US]; 116 Hunter Court, Wilmington, DE 19808 (US).STERNBERG, Charlene, Gross [US/US]; 3111 Loudoun
Drive, Wilmington, DE 19808-2703 (US).(74) Agents: MAYER, Nancy, S. et al.; E.I. du Pont de Nemours
and Company, Legal/Patent Records Center, 11007 Market
Street, Wilmington, DE 19898 (US).(81) Designated States: AU, BB, BG, BR, BY, CA, CN, CZ, FI,
GE, HU, JP, KG, KP, KR, KZ, LK, LV, MD, MG, MN,
MW, NO, NZ, PL, RO, RU, SD, SI, SK, TJ, TT, UA, US,
UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF,
BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).


Published

With international search report.
With amended claims.


(54) Title: FUNGICIDAL FUSED BICYCLIC PYRIMIDINONES



(I)



(II)



(III)

(57) Abstract

This invention pertains to compounds of Formulae (I), (II), and (III) including all geometric and stereoisomers, *N*-oxides, agriculturally-suitable salts thereof, agricultural compositions containing them and their use as fungicides, wherein: Q is O or S; and n, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, and R⁹ are described in the text.

5117-P1-41-EMA

-USSN 08/545,241

Filed November 21, 1995

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

TITLE




FUNGICIDAL FUSED BICYCLIC PYRIMIDINONES

This invention relates to certain 4(3H)-quinazolinones, their agriculturally suitable salts and compositions, and methods of their use as general or selective fungicides, in particular for the control of cereal powdery mildew both preventive and curative.

U.S. 3,755,582 and U.S. 3,867,384 disclose certain 4(3H)-quinazolinone fungicides. These patents, however, do not specifically disclose the compounds of the present invention.

SUMMARY OF THE INVENTION

This invention comprises compounds of Formulae I, II, and III including all geometric and stereoisomers, *N*-oxides, agriculturally-suitable salts thereof, agricultural compositions containing them and their use as fungicides:



15

wherein:

n is 0, 1 or 2;

Q is independently O or S;

*R*¹ is C₃-C₁₀ alkyl; C₃-C₅ cycloalkyl; C₄-C₁₀ alkenyl; C₄-C₁₀ alkynyl; C₁-C₁₀ haloalkyl; C₃-C₁₀ haloalkenyl; C₃-C₁₀ haloalkynyl; C₂-C₁₀ alkoxyalkyl; C₂-C₁₀ alkylthioalkyl; C₂-C₁₀ alkylsulfinylalkyl; C₂-C₁₀ alkylsulfonylalkyl; C₅-C₁₀ cycloalkylalkyl; C₄-C₁₀ alkenyloxyalkyl; C₄-C₁₀ alkynyloxyalkyl; C₄-C₁₀ (cycloalkyl)oxyalkyl; C₄-C₁₀ alkenylthioalkyl; C₄-C₁₀ alkynylthioalkyl; C₆-C₁₀ (cycloalkyl)thioalkyl; C₂-C₁₀ haloalkoxyalkyl; C₄-C₁₀ haloalkenyloxyalkyl; C₄-C₁₀ haloalkynyloxyalkyl; C₄-C₁₀ alkoxyalkenyl; C₄-C₁₀ alkoxyalkynyl; C₄-C₁₀ alkylthioalkenyl; C₄-C₁₀ alkylthioalkynyl; C₄-C₁₀ trialkylsilylalkyl; C₁-C₁₀ alkyl substituted with NR¹¹R¹², nitro, cyano, or phenyl optionally substituted with R¹⁴, R¹⁵, and R¹⁶; C₁-C₁₀ alkoxy; C₁-C₁₀ haloalkoxy; C₁-C₁₀ alkylthio; C₁-C₁₀ haloalkylthio; NR¹¹R¹²; or pyridyl, furanyl, thienyl, naphthyl, benzofuranyl, benzothienyl, or quinolinyl each optionally substituted with R¹⁴, R¹⁵, and R¹⁶;

- R^2 is C_3 - C_{10} alkyl; C_6 - C_7 cycloalkyl; C_3 - C_{10} alkenyl; C_3 - C_{10} alkynyl; C_1 - C_{10} haloalkyl; C_3 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_2 - C_{10} alkoxyalkyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_4 - C_{10} cycloalkylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_3 - C_{10} cyanoalkyl; C_2 - C_{10} nitroalkyl; C_1 - C_{10} alkyl substituted with CO_2R^{11} , $NR^{11}R^{12}$, or phenyl optionally substituted with R^{13} , R^{15} , and R^{16} ; phenyl optionally substituted with R^{13} , R^{15} , and R^{16} ; $-N=CR^{11}R^{11}$; or $-NR^{11}R^{12}$; or
- R^1 and R^2 are taken together to form $-CH_2(CH_2)_mCH_2-$; m is 1-4;
- R^3 is halogen; C_1 - C_8 alkyl; C_3 - C_8 cycloalkyl; C_2 - C_8 alkenyl; C_2 - C_8 alkynyl; C_1 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; C_3 - C_8 haloalkynyl; C_1 - C_8 alkoxy; C_1 - C_8 haloalkoxy; C_3 - C_8 alkenyloxy; C_3 - C_8 alkynyloxy; C_1 - C_8 alkylthio; C_3 - C_8 alkenylthio; C_3 - C_8 alkynylthio; C_1 - C_8 alkylsulfinyl; C_1 - C_8 alkylsulfonyl; C_2 - C_8 alkoxyalkyl; C_2 - C_8 alkylthioalkyl; C_2 - C_8 alkylsulfinylalkyl; C_2 - C_8 alkylsulfonylalkyl; C_4 - C_8 cycloalkylalkyl; C_3 - C_8 trialkylsilyl; nitro; $NR^{11}R^{12}$; C_5 - C_8 trialkylsilylalkynyl; or phenyl optionally substituted with at least one R^{13} ;
- R^4 is hydrogen; halogen; C_1 - C_4 alkyl; C_1 - C_4 haloalkyl; C_1 - C_4 alkoxy; or C_1 - C_4 haloalkoxy;
- R^5 is C_3 - C_5 alkyl; C_7 - C_{10} alkyl; C_4 - C_7 alkenyl; C_3 - C_5 alkynyl; C_1 - C_{10} haloalkyl; C_5 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_2 - C_{10} alkoxyalkyl other than methoxypropyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_4 - C_{10} cycloalkylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_1 - C_{10} alkyl substituted with $NR^{11}R^{12}$, nitro, or phenyl optionally substituted with at least one of R^{14} , R^{15} , and R^{16} ; C_2 - C_{10} alkyl substituted with cyano; C_1 - C_{10} alkoxy; C_1 - C_{10} haloalkoxy; C_1 - C_{10} alkylthio; C_1 - C_{10} haloalkylthio; $NR^{11}R^{12}$; or phenyl, furanyl, thienyl, naphthyl, benzofuranyl, or benzothienyl each optionally substituted with R^{14} , R^{15} , and R^{16} ;

5
10
15
20
25
30
35

R^6 is C_3 - C_{10} alkyl; C_3 - C_7 alkenyl; C_3 - C_{10} alkynyl; C_1 - C_{10} haloalkyl; C_3 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_3 - C_{10} alkoxyalkyl other than propoxymethyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_4 - C_{10} cycloalkylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_5 - C_{10} cyanoalkyl; C_2 - C_{10} nitroalkyl; or C_3 - C_{10} alkyl substituted with CO_2R^{11} , $NR^{11}R^{12}$, or phenyl optionally substituted with R^{13} , R^{15} , and R^{16} ; or phenyl optionally substituted with R^{13} , R^{15} , and R^{16} ; or

R^5 and R^6 are taken together to form $-CH_2(CH_2)_mCH_2-$;

15
20
25
30
35

R^7 is C_3 - C_{10} alkyl; C_3 - C_7 cycloalkyl; C_4 - C_7 alkenyl; propynyl; C_5 - C_{10} alkynyl; C_2 - C_{10} haloalkyl; C_3 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_2 - C_{10} alkoxyalkyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_1 - C_{10} alkyl substituted with $NR^{11}R^{12}$ or nitro; C_2 - C_{10} alkyl substituted with cyano; C_1 - C_{10} alkoxy; C_1 - C_{10} haloalkoxy; C_1 - C_{10} alkylthio; C_1 - C_{10} haloalkylthio; $NR^{12}R^{17}$; or phenyl, pyridyl, furanyl, thienyl, naphthyl, benzofuranyl, benzothienyl, or quinolinyl each optionally substituted with R^{14} , R^{15} , and R^{16} ;

R^8 is hydrogen; C_1 - C_4 alkyl; or $-C(=O)R^{10}$;

30
35

R^9 is hydrogen; C_2 - C_{10} alkyl; C_3 - C_7 cycloalkyl; C_3 - C_{10} alkenyl; C_3 - C_{10} alkynyl; C_3 - C_{10} haloalkyl; C_3 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_3 - C_{10} alkoxyalkyl other than butoxyethyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_4 - C_{10} cycloalkylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_1 - C_{10} alkyl substituted with $NR^{11}R^{12}$; C_4 - C_{10} cyanoalkyl; C_2 - C_{10} nitroalkyl; C_1 - C_8 alkyl substituted with CO_2R^{11} ; pyridyl, furanyl, thienyl, or

- naphthyl each optionally substituted with R^{14} , R^{15} , and R^{16} ; $-N=CR^{11}R^{11}$; $-NR^{12}R^{17}$; $-OR^{12}$; or $-NC(=Q)NR^{11}R^{12}$; or R^3 and R^4 are both iodine and R^9 is phenyl optionally substituted with R^{14} , R^{15} , and R^{16} ; or
- R^7 and R^9 are taken together to form $-\text{CH}_2(\text{CH}_2)_m\text{CH}_2-$;
- R^{10} is hydrogen; C_1 - C_4 alkyl; C_1 - C_4 alkoxy; or $NR^{11}R^{12}$;
- R^{11} is independently hydrogen; C_1 - C_4 alkyl; or phenyl optionally substituted with at least one R^{13} ;
- R^{12} is independently hydrogen; C_1 - C_8 alkyl; or phenyl optionally substituted with at least one R^{13} ; or
- R^{11} and R^{12} are taken together to form $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2(\text{CH}_2)_3\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}(\text{Me})\text{CH}_2\text{CH}(\text{Me})\text{CH}_2-$, or $-\text{CH}_2\text{CH}(\text{Me})\text{OCH}(\text{Me})\text{CH}_2-$;
- R^{13} is independently halogen; C_1 - C_4 alkyl; C_1 - C_4 alkoxy; C_1 - C_4 haloalkyl; nitro; or cyano;
- R^{14} is independently C_1 - C_6 alkyl; C_1 - C_6 alkoxy; C_1 - C_6 haloalkyl; halogen; C_2 - C_8 alkynyl; C_1 - C_6 thioalkyl; phenyl or phenoxy each optionally substituted with at least one R^{13} ; cyano; nitro; C_1 - C_6 haloalkoxy; C_1 - C_6 haloalkylthio; C_2 - C_6 alkenyl; C_2 - C_6 haloalkenyl; acetyl; CO_2Me ; or $\text{N}(\text{C}_1\text{-C}_2 \text{ alkyl})_2$;
- R^{15} is independently methyl; ethyl; methoxy; methylthio; halogen; or trifluoromethyl;
- R^{16} is independently halogen; and
- R^{17} is independently C_1 - C_8 alkyl; or phenyl optionally substituted with at least one R^{13} .

DETAILED DESCRIPTION OF THE INVENTION

- In the above recitations, the term "alkyl", used either alone or in compound words such as "alkylthio," "haloalkyl," or "alkylthioalkyl" denotes straight-chain or branched alkyl; e.g., methyl, ethyl, *n*-propyl, *i*-propyl, or the different butyl, pentyl, hexyl, etc. isomers.

- "Cycloalkyl" denotes cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl. The term "cycloalkyloxyalkyl" denotes the cycloalkyl groups linked through an oxygen atom to an alkyl chain. Examples include cyclopentyloxymethyl and cyclohexyloxybutyl. The term "cycloalkylthioalkyl" are the cycloalkyl groups linked through a sulfur atom to an alkyl chain; e.g., cyclopropylthiopentyl. "Cycloalkylalkyl" denotes a cycloalkyl ring attached to a branched or straight-chain alkyl; e.g. cyclopropylmethyl and cyclohexylbutyl.

"Alkenyl" denotes straight chain or branched alkenes; e.g., 1-propenyl, 2-propenyl, 3-propenyl and the different butenyl, pentenyl, hexenyl, etc. isomers. Alkenyl also denotes polyenes such as 1,3-hexadiene and 2,4,6-heptatriene.

"Alkynyl" denotes straight chain or branched alkynes; e.g., ethynyl, 1-propynyl, 3-propynyl and the different butynyl, pentynyl, hexynyl, etc. isomers. "Alkynyl" can also denote moieties comprised of multiple triple bonds; e.g., 2,7-octadiyne and 2,5,8-decatriyne.

- 5 "Alkoxy" denotes methoxy, ethoxy, *n*-propyloxy, isopropyloxy and the different butoxy, pentoxy, hexyloxy, etc. isomers. "Alkoxyalkenyl" and "alkoxyalkynyl" denoted groups in which the alkoxy group is bonded through the oxygen atom to an alkenyl or alkynyl group, respectively. Examples include $\text{CH}_3\text{OCH}_2\text{CH}=\text{CH}$ and $(\text{CH}_3)_2\text{CHOCH}_2\text{C}\equiv\text{CCH}_2$. The corresponding sulfur derivatives are denoted
- 10 "alkylthioalkenyl and "alkylthioalkynyl." Examples of the former include $\text{CH}_3\text{SCH}_2\text{CH}=\text{CH}$ and $\text{CH}_3\text{CH}_2\text{SCH}_2(\text{CH}_3)\text{CH}=\text{CHCH}_2$, and an example of the latter is $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{SCH}_2\text{C}\equiv\text{C}$.

- "Alkenyloxy" denotes straight chain or branched alkenyloxy moieties. Examples of alkenyloxy include $\text{H}_2\text{C}=\text{CHCH}_2\text{O}$, $(\text{CH}_3)_2\text{C}=\text{CHCH}_2\text{O}$, $(\text{CH}_3)\text{CH}=\text{CHCH}_2\text{O}$, $(\text{CH}_3)\text{CH}=\text{C}(\text{CH}_3)\text{CH}_2\text{O}$ and $\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{O}$. "Alkenylthio" denotes the similar groups wherein the oxygen atom is replaced with a sulfur atom; e.g., $\text{H}_2\text{C}=\text{CHCH}_2\text{S}$ and $(\text{CH}_3)\text{CH}=\text{C}(\text{CH}_3)\text{CH}_2\text{S}$. The term "alkenyloxyalkyl" denotes groups in which the alkenyloxy moiety is attached to an alkyl group. Examples include
- 15 $\text{H}_2\text{C}=\text{CHCH}_2\text{OCH}_2\text{CH}_2$, $\text{H}_2\text{C}=\text{CHCH}_2\text{OCH}(\text{CH}_3)\text{CH}_2$, etc. "Alkenylthioalkyl" denotes the alkenylthio moieties bonded to an alkyl group. Examples include $\text{H}_2\text{C}=\text{CHCH}_2\text{SCH}(\text{CH}_3)\text{CH}(\text{CH}_3)$ and $(\text{CH}_3)\text{CH}=\text{C}(\text{CH}_3)\text{CH}_2\text{SCH}_2$.

- "Alkynyloxy" denotes straight or branched alkynyloxy moieties. Examples include $\text{HC}\equiv\text{CCH}_2\text{O}$, $\text{CH}_3\text{C}\equiv\text{CCH}_2\text{O}$ and $\text{CH}_3\text{C}\equiv\text{CCH}_2\text{CH}_2\text{O}$. "Alkynyloxyalkyl" denotes alkynyloxy moieties bonded to alkyl groups; e.g., $\text{CH}_3\text{C}\equiv\text{CCH}_2\text{OCH}_2\text{CH}_2$ and
- 25 $\text{HC}\equiv\text{CCH}_2\text{OCH}(\text{CH}_3)\text{CH}_2$. "Alkynylthioalkyl" denotes alkynylthio moieties bonded to alkyl groups. Example include $\text{CH}_3\text{C}\equiv\text{CCH}_2\text{SCH}_2\text{CH}_2$ and $\text{CH}_3\text{C}\equiv\text{CCH}_2\text{CH}_2\text{SCH}(\text{CH}_3)\text{CH}_2$.

- "Alkylthio" denotes methylthio, ethylthio, and the different propylthio, butylthio, pentylthio and hexylthio isomers. "Alkylthioalkyl" denotes alkylthio groups attached to
- 30 an alkyl chain; e.g., $\text{CH}_3\text{CH}_2\text{SCH}_2\text{CH}(\text{CH}_3)$ and $(\text{CH}_3)_2\text{CHSCH}_2$.

- "Alkylsulfinyl" denotes both enantiomers of an alkylsulfinyl group. For example, $\text{CH}_3\text{S}(\text{O})$, $\text{CH}_3\text{CH}_2\text{S}(\text{O})$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{S}(\text{O})$, $(\text{CH}_3)_2\text{CHS}(\text{O})$ and the different butylsulfinyl, pentylsulfinyl and hexylsulfinyl isomers. "Alkylsulfinylalkyl" denotes alkylsulfinyl groups attached to an alkyl chain; e.g., $\text{CH}_3\text{CH}_2\text{S}(\text{O})\text{CH}_2\text{CH}(\text{CH}_3)$ and
- 35 $(\text{CH}_3)_2\text{CHS}(\text{O})\text{CH}_2$.

Examples of "alkylsulfonyl" include $\text{CH}_3\text{S}(\text{O})_2$, $\text{CH}_3\text{CH}_2\text{S}(\text{O})_2$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{S}(\text{O})_2$, $(\text{CH}_3)_2\text{CHS}(\text{O})_2$ and the different butylsulfonyl, pentylsulfonyl and

hexylsulfonyl isomers. "Alkylsulfonylalkyl" denotes alkylsulfonyl groups attached to an alkyl chain; e.g., $\text{CH}_3\text{CH}_2\text{S}(\text{O})_2\text{CH}_2\text{CH}(\text{CH}_3)$ and $(\text{CH}_3)_2\text{CHS}(\text{O})_2\text{CH}_2$.

The term "halogen", either alone or in compound words such as "haloalkyl", denotes fluorine, chlorine, bromine or iodine. Further, when used in compound words such as "haloalkyl", said alkyl may be partially or fully substituted with halogen atoms which may be the same or different. Examples of "haloalkyl" include F_3C , ClCH_2 , CF_3CH_2 and CF_3CF_2 . Examples of "haloalkenyl" include $(\text{Cl})_2\text{C}=\text{CHCH}_2$ and $\text{CF}_3\text{CH}_2\text{CH}=\text{CHCH}_2$. "Haloalkenyloxyalkyl" denotes haloalkenyl groups bonded to oxygen and in turn bonded to alkyl groups. Examples include $\text{CF}_3\text{CH}_2\text{CH}=\text{CHCH}_2\text{OCH}_2$ and $(\text{Cl})_2\text{C}=\text{CHCH}_2\text{OCH}_2\text{CH}_2$. Examples of "haloalkynyl" include $\text{HC}\equiv\text{CCHCl}$, $\text{CF}_3\text{C}\equiv\text{C}$, $\text{CCl}_3\text{C}\equiv\text{C}$ and $\text{FCH}_2\text{C}\equiv\text{CCH}_2$. "Haloalkynyloxyalkyl" denotes haloalkynyl groups bonded through an oxygen atom to an alkyl moiety. Examples include $\text{CF}_3\text{C}\equiv\text{CCH}_2\text{OCH}_2\text{CH}_2$, $\text{ClCH}_2\text{C}\equiv\text{CCH}_2\text{CH}_2\text{OCH}(\text{CH}_3)$, etc. Examples of "haloalkoxy" include CF_3O , $\text{CCl}_3\text{CH}_2\text{O}$, $\text{CF}_2\text{HCH}_2\text{CH}_2\text{O}$ and $\text{CF}_3\text{CH}_2\text{O}$. "Haloalkoxyalkyl" denotes haloalkoxy groups bonded to straight-chain or branched alkyl groups; e.g., $\text{CF}_2\text{HCH}_2\text{CH}_2\text{OCH}_2\text{CH}_2$, $\text{CCl}_3\text{CH}_2\text{OCH}(\text{CH}_3)$ and CF_3OCH_2 .

"Trialkylsilyl" designates a group with three alkyl groups bonded to silicon; e.g., $(\text{CH}_3)_3\text{Si}$ and $t\text{-Bu}(\text{CH}_3)_2\text{Si}$. "Trialkylsilylalkyl" denotes trialkylsilyl groups bonded to another straight-chain or branched alkyl group. Examples include $(\text{CH}_3)_3\text{SiCH}_2$ and $t\text{-Bu}(\text{CH}_3)_2\text{SiCH}_2\text{CH}(\text{CH}_3)\text{CH}_2$.

The total number of carbon atoms in a substituent group is indicated by the " $\text{C}_i\text{-C}_j$ " prefix where i and j are numbers from 1 to 10. For example, $\text{C}_1\text{-C}_3$ alkylsulfonyl designates methylsulfonyl through propylsulfonyl; C_2 alkoxyalkoxy designates $\text{CH}_3\text{OCH}_2\text{O}$; C_3 alkoxyalkoxy designates, for example, $\text{CH}_3\text{OCH}_2\text{CH}_2\text{O}$ or $\text{CH}_3\text{CH}_2\text{OCH}_2\text{O}$; and C_4 alkoxyalkoxy designates the various isomers of an alkoxy group substituted with a second alkoxy group containing a total of 4 carbon atoms, examples including $\text{CH}_3\text{CH}_2\text{CH}_2\text{OCH}_2\text{O}$, and $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_2\text{O}$. Examples of "alkoxyalkyl" include CH_3OCH_2 , $\text{CH}_3\text{OCH}_2\text{CH}_2$, $\text{CH}_3\text{CH}_2\text{OCH}_2$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{OCH}_2$ and $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_2$.

Preferred for reasons of ease of synthesis or greater fungicidal activity are:

Preferred 1. The compounds of Formula I as defined above wherein:

Q is O;

R^1 is $\text{C}_3\text{-C}_8$ alkyl; $\text{C}_4\text{-C}_8$ alkenyl; $\text{C}_4\text{-C}_8$ alkynyl; $\text{C}_1\text{-C}_8$ haloalkyl; $\text{C}_3\text{-C}_8$ haloalkenyl; $\text{C}_2\text{-C}_8$ alkoxyalkyl; $\text{C}_2\text{-C}_8$ alkylthioalkyl; $\text{C}_5\text{-C}_8$ cycloalkylalkyl; $\text{C}_2\text{-C}_8$ alkyl substituted with cyano; $\text{C}_1\text{-C}_8$ alkoxy; $\text{C}_1\text{-C}_8$ haloalkoxy; $\text{C}_1\text{-C}_8$ alkylthio; or $\text{C}_4\text{-C}_8$ alkenyloxyalkyl; or pyridyl, furanyl, or thienyl each optionally substituted with R^{14} and R^{15} ;

R² is C₃-C₈ alkyl; C₃-C₈ alkenyl; C₃-C₈ alkynyl; C₁-C₈ haloalkyl; C₃-C₈ haloalkenyl; C₃-C₈ alkoxyalkyl; C₂-C₈ alkylthioalkyl; C₄-C₈ cycloalkylalkyl; C₃-C₈ cyanoalkyl; C₄-C₈ alkenyloxyalkyl; or phenyl optionally substituted with R¹³;

5 R³ is halogen; C₁-C₈ alkyl; C₂-C₈ alkynyl; C₃-C₈ cycloalkyl; C₁-C₈ haloalkyl; C₁-C₈ alkoxy; C₁-C₈ haloalkoxy; C₁-C₈ alkylthio; C₁-C₈ alkylsulfonyl; C₂-C₈ alkoxyalkyl; C₂-C₈ alkylthioalkyl; C₄-C₈ cycloalkylalkyl; or C₅-C₈ trialkylsilylalkynyl; and

10 R¹⁴ is methyl; ethyl; methoxy; ethoxy; C₁-C₂ haloalkyl; halogen; acetylenyl; propargyl; methylthio; ethylthio; cyano; nitro; C₁-C₂ haloalkoxy; vinyl; allyl; acetyl; CO₂Me; or N(C₁-C₂ alkyl)₂.

Preferred 2. The compounds of Formula II as defined above wherein:

Q is O;

n is 0;

15 R³ is halogen; C₁-C₈ alkyl; C₂-C₈ alkynyl; C₃-C₈ cycloalkyl; C₁-C₈ haloalkyl; C₁-C₈ alkoxy; C₁-C₈ haloalkoxy; C₁-C₈ alkylthio; C₁-C₈ alkylsulfonyl; C₂-C₈ alkoxyalkyl; C₂-C₈ alkylthioalkyl; C₄-C₈ cycloalkylalkyl; or C₅-C₈ trialkylsilylalkynyl;

20 R⁵ is C₃-C₅ alkyl; C₄-C₇ alkenyl; C₃-C₅ alkynyl; C₁-C₈ haloalkyl; C₅-C₈ haloalkenyl; C₂-C₈ alkoxyalkyl other than methoxypropyl; C₂-C₈ alkylthioalkyl; C₄-C₈ cycloalkylalkyl; C₂-C₈ alkyl substituted with cyano; C₁-C₈ alkoxy; C₁-C₈ haloalkoxy; C₁-C₈ alkylthio; or C₄-C₈ alkenyloxyalkyl; or phenyl, furanyl, or thienyl each optionally substituted with R¹⁴ and R¹⁵;

25 R⁶ is C₃-C₈ alkyl; C₃-C₇ alkenyl; C₃-C₈ alkynyl; C₁-C₈ haloalkyl; C₃-C₈ haloalkenyl; C₃-C₈ alkoxyalkyl other than propoxymethyl; C₂-C₈ alkylthioalkyl; C₄-C₈ cycloalkylalkyl; C₅-C₈ cyanoalkyl; C₄-C₈ alkenyloxyalkyl; phenyl optionally substituted with R¹³; or C₃-C₅ alkyl substituted with phenyl optionally substituted with R¹³ and R¹⁵; and

30 R¹⁴ is methyl; ethyl; methoxy; ethoxy; C₁-C₂ haloalkyl; halogen; acetylenyl; propargyl; methylthio; ethylthio; cyano; nitro; C₁-C₂ haloalkoxy; vinyl; allyl; acetyl; CO₂Me; or N(C₁-C₂ alkyl)₂.

Preferred 3. The compounds of Formula III as defined above wherein:

35 Q is O;

R³ is halogen; C₁-C₈ alkyl; C₂-C₈ alkynyl; C₃-C₈ cycloalkyl; C₁-C₈ haloalkyl; C₁-C₈ alkoxy; C₁-C₈ haloalkoxy; C₁-C₈ alkylthio; C₁-C₈

alkylsulfonyl; C₂-C₈ alkoxyalkyl; C₂-C₈ alkylthioalkyl; C₄-C₈ cycloalkylalkyl; or C₅-C₈ trialkylsilylalkynyl;

R⁷ is C₃-C₈ alkyl; C₄-C₇ alkenyl; propynyl; C₂-C₈ haloalkyl; C₃-C₈ haloalkenyl; C₂-C₈ alkoxyalkyl; C₂-C₈ alkylthioalkyl; C₂-C₈ alkyl substituted with cyano; C₁-C₈ alkoxy; C₁-C₈ haloalkoxy; C₁-C₈ alkylthio; or C₄-C₈ alkenyloxyalkyl; or phenyl, pyridyl, furanyl, or thienyl each optionally substituted with R¹⁴ and R¹⁵;

R⁹ is C₃-C₈ alkyl; C₃-C₈ alkenyl; C₃-C₈ alkynyl; C₃-C₈ haloalkyl; C₃-C₈ haloalkenyl; C₃-C₈ alkoxyalkyl; C₂-C₈ alkylthioalkyl; C₄-C₈ cycloalkylalkyl; C₄-C₈ cyanoalkyl; C₄-C₈ alkenyloxyalkyl; -NR¹²R¹⁷; or R³ and R⁴ are both iodine and R⁹ is phenyl optionally substituted with R¹⁴ and R¹⁵; and

R¹⁴ is methyl; ethyl; methoxy; ethoxy; C₁-C₂ haloalkyl; halogen; acetylenyl; propargyl; methylthio; ethylthio; cyano; nitro; C₁-C₂ haloalkoxy; vinyl; allyl; acetyl; CO₂Me; or N(C₁-C₂ alkyl)₂.

Preferred 4. The compounds of Preferreds 1, 2, and 3 wherein:

R¹ is C₃-C₈ alkyl; C₄-C₈ alkenyl; C₄-C₈ alkynyl; C₃-C₈ haloalkyl; C₃-C₈ haloalkenyl; C₃-C₈ alkoxyalkyl; or thienyl optionally substituted with at least one of R¹⁴ and R¹⁵;

R² is C₃-C₈ alkyl; C₃-C₈ alkenyl; C₃-C₈ alkynyl; C₃-C₈ haloalkyl; C₃-C₈ haloalkenyl; C₃-C₈ alkoxyalkyl; or phenyl optionally substituted with R¹³;

R³ is halogen; C₁-C₄ alkyl; C₁-C₄ haloalkyl; C₁-C₄ alkoxy; C₁-C₄ haloalkoxy; acetylenyl; or trimethylsilylacetylenyl;

R⁵ is C₃-C₅ alkyl; C₄-C₇ alkenyl; C₃-C₅ alkynyl; C₃-C₈ haloalkyl; C₅-C₈ haloalkenyl; C₃-C₈ alkoxyalkyl; or phenyl or thienyl each optionally substituted with R¹⁴ and R¹⁵;

R⁶ is C₃-C₈ alkyl; C₃-C₇ alkenyl; C₃-C₈ alkynyl; C₃-C₈ haloalkyl; C₃-C₈ haloalkenyl; C₃-C₈ alkoxyalkyl; or phenyl optionally substituted with R¹³;

R⁷ is C₃-C₈ alkyl; C₄-C₇ alkenyl; propynyl; C₃-C₈ haloalkyl; C₃-C₈ haloalkenyl; C₃-C₈ alkoxy; C₃-C₈ alkoxyalkyl; or phenyl or thienyl each optionally substituted with R¹⁴ and R¹⁵;

R⁹ is C₃-C₈ alkyl; C₃-C₈ alkenyl; C₃-C₈ alkynyl; C₃-C₈ haloalkyl; C₃-C₈ haloalkenyl; C₃-C₈ alkoxyalkyl; -NR¹²R¹⁷; or R³ and R⁴ are both iodine and R⁹ is phenyl optionally substituted with R¹⁴ and R¹⁵; and

R¹⁴ is methyl; ethyl; methoxy; methylthio; halogen; trifluoromethyl; or N(C₁-C₂ alkyl)₂.

Preferred 5. The compounds of Preferred 4 wherein:

R^1 is C_3 - C_8 alkyl; C_4 - C_8 alkenyl; C_4 - C_8 alkynyl; C_3 - C_8 haloalkyl; or C_3 - C_8 haloalkenyl;

R^2 is C_3 - C_8 alkyl; C_3 - C_8 alkenyl; C_3 - C_8 alkynyl; C_3 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; or phenyl optionally substituted with R^{13} ;

R^3 is halogen;

R^4 is hydrogen or halogen;

R^5 is C_3 - C_5 alkyl; C_4 - C_7 alkenyl; C_3 - C_5 alkynyl; C_3 - C_8 haloalkyl; or C_5 - C_8 haloalkenyl; or phenyl optionally substituted with R^{14} and R^{15} ;

R^6 is C_3 - C_8 alkyl; C_3 - C_7 alkenyl; C_3 - C_8 alkynyl; C_3 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; or phenyl optionally substituted with R^{13} ;

R^7 is C_3 - C_8 alkyl; C_4 - C_7 alkenyl; propynyl; C_3 - C_8 haloalkyl; or C_3 - C_8 haloalkenyl; or phenyl optionally substituted with R^{14} and R^{15} ; and

R^9 is C_3 - C_8 alkyl; C_3 - C_8 alkenyl; C_3 - C_8 alkynyl; C_3 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; $-NR^{12}R^{17}$; or R^3 and R^4 are both iodine and R^9 is phenyl optionally substituted with R^{14} and R^{15} .

Preferred 6. The compounds of Preferred 5 wherein said compounds are selected from the group:

6-bromo-3-propyl-2-propyloxy-4(3H)-quinazolinone;
6,8-diiodo-3-propyl-2-propyloxy-4(3H)-quinazolinone;
6-iodo-3-propyl-2-propyloxy-4(3H)-quinazolinone; and
6,8-diiodo-3-propyl-2-(phenylamino)-4(3H)-quinazolinone.

It is recognized that some reagents and reaction conditions described below for preparing compounds of Formulae I, II, and III may not be compatible with some functionalities claimed for R^1 - R^{17} , n, m, and Q. In these cases, the incorporation of protection/deprotection sequences into the synthesis may be necessary in order to obtain the desired products. The cases in which protecting groups are necessary, and which protecting group to use, will be apparent to one skilled in chemical synthesis. See Greene, T. W. and Wuts, P. G. M.; *Protective Groups in Organic Synthesis*, 2nd Ed.; John Wiley & Sons, Inc.; New York, (1980) for suitable protecting groups.

In the following description of the preparation of compounds of Formulae I, II, and III, compounds of Formulae Ia and Ib, IIa-IIc, and IIIa-IIIe are various subsets of the compounds of Formulae I, II, and III. All substituents in compounds of Formulae Ia and Ib, IIa-IIc, and IIIa-IIIe and 2-7 are as defined above for Formulae I, II, and III.

Compounds of this invention can exist as one or more stereoisomers. The various stereoisomers include enantiomers, diastereomers and geometric isomers. One skilled in

the art will appreciate that one stereoisomer may be more active than the others and how to separate said stereoisomers. Accordingly, the present invention comprises mixtures, individual stereoisomers, and optically active mixtures of compounds of Formulae I, II, and III as well as agriculturally suitable salts thereof.

- 5 The compounds of Formulae I, II, and III can be prepared as described below in Schemes 1-9 and Examples 1-3.


Synthesis of Compounds of Formula I

Compounds of Formula Ia, compounds of Formula I wherein Q is O, can be made by the method illustrated in Scheme 1.

- 10 An anthranilic acid (2-aminobenzoic acid) of Formula 2 is condensed with an isothiocyanate of Formula $R^1\text{-NCS}$ to form the 2-thioquinazolinone of Formula 3. This condensation is preferably performed in the presence of a base such as triethylamine. S-Methylation of this compound affords the 2-methylthio-4(3H)-quinazolinone of Formula 4.


- 15 For the introduction of the $R^2\text{O}$ group, the 2-methylthio-4(3H)-quinazolinone of Formula 4 is treated with a mixture of a base, for example sodium hydride, in $R^2\text{OH}$ solvent. The reaction mixture is stirred at a temperature from about 0°C to 120°C for 1-120 hours. The desired 2- $R^2\text{O}$ -4(3H)-quinazolinone can be isolated from the reaction mixture by extraction into a water-immiscible solvent, and purified by chromatography or
20 recrystallization. Similar synthetic procedures are described in U.S. 3,755,582, incorporated herein by reference.

11

Scheme 1

- Anthranilic acids of Formula 2 are known or can be prepared by known methods. For example see, March, J. *Advanced Organic Chemistry*; 3rd ed., John Wiley: New York, (1985), p 983. The isothiocyanates of Formula R¹-NCS can be prepared from the corresponding amine by treatment with thiophosgene as known in the art. For example, see *J. Heterocycl. Chem.*, (1990), 27, 407.


- Alternatively, 2-thioquinazolin-4(1H)-ones of Formula 3 can be prepared by treatment of the (C₁-C₄ alkyl) anthranilic acid ester of Formula 5 with thiophosgene to form the isothiocyanate ester, followed by treatment with an amine of formula R¹NH₂ (Scheme 2).

Scheme 2

The anthranilic acid ester of Formula 5 is treated with thiophosgene at a temperature from about -20°C to 100°C for 1 to 48 hours optionally in an inert solvent. Often this reaction is performed in a biphasic mixture in the presence of a base, such as calcium carbonate, and an acid, such as aqueous hydrochloric acid. The resulting isothiocyanate may be isolated by extraction into a water-immiscible solvent, such as methylene chloride, followed by drying of the organic extracts and evaporation under reduced pressure. Alternatively, the isothiocyanate can be combined in situ with the amine of Formula H_2NR^1 and stirred at about -20°C to 50°C for 0.1 to 24 hours. The desired 2-thioquinazolin-4(3H)-ones of Formula 3 can be isolated from the reaction mixture by aqueous extraction, and purified by chromatography or recrystallization. Similar synthetic procedures are described in *J. Heterocycl. Chem.*, (1990), 27, 407.

Compounds of Formula Ib, compounds of Formula I wherein Q is S, can be prepared as illustrated in Scheme 3.

Scheme 3



15

Treatment of the quinazolinone of Formula Ia with phosphorous pentasulfide or Lawesson's reagent [2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulfide] in an inert solvent such as dioxane at a temperature from 0°C to the reflux temperature of the solvent for 0.1 to 72 hours affords the quinazolinethione of Formula Ib. This procedure is described in the literature, for example see U.S. 3,755,582.


20

Synthesis of Compounds of Formula II

4(3H)-Quinazolinones of Formula IIa, compounds of Formula II wherein n is 0 and Q is O, can be prepared by a modification of the synthesis illustrated in Scheme 1. As illustrated in Scheme 4, the 2-thioquinazolin-4(3H)-one of Formula 6 is alkylated with R^6-X wherein X is a typical leaving group such as Br, I, CH_3SO_3 (OMs), or $(4-CH_3-Ph)SO_3$ (OTs) to afford the 2- R^6 S-4(3H)-quinazolinone of Formula IIa. One or more equivalents of a base can be used to accelerate the process. Bases such as sodium hydroxide and sodium hydride are suitable.

30


13

Scheme 4

Typically, the 2-thioquinazolin-4(3H)-one is dissolved or dispersed in an inert solvent such as dimethylformamide and treated with a base at a temperature from about -20°C to 60°C with a base. The reaction mixture may then be heated to just above ambient temperature to the reflux temperature of the solvent for 0.1 to 24 hours to effect deprotonation. After cooling, the reaction mixture is cooled and treated with R⁶-X and stirred for 0.1-24 hours at a temperature from about 20°C to the reflux temperature of the solvent. The quinazolinone of Formula IIa can be isolated by extraction into a water-immiscible solvent, and purified by chromatography or recrystallization.

2-Thioquinazolin-4(3H)-ones of Formula 6 are prepared as described above in Schemes 1 and 2 for 2-thioquinazolin-4(3H)-ones of Formula 3.


4(3H)-Quinazolinones of Formula IIb, compounds of Formula II wherein Q is O and n is 1 or 2, can be prepared by oxidation of the corresponding -SR⁶ compound of Formula Ia using well-known procedures for oxidation of sulfur (Scheme 5). For example, see March, J. *Advanced Organic Chemistry*; 3rd ed., John Wiley: New York, (1985), p 1089.

Scheme 5


20

4(3H)-Quinazolinethiones of Formula IIc, compounds of Formula II wherein Q is S, can be prepared by treatment of the corresponding quinazolinone with phosphorous pentasulfide or Lawesson's reagent as described in U.S. 3,755,582 and above for compounds of Formula Ib (Scheme 6).

14

Scheme 6Synthesis of Compounds of Formula III

- 5 4(3H)-Quinazolinones of Formula IIIa, compounds of Formula III wherein Q is O, can be prepared by the method illustrated in Scheme 7. This method is described in detail in U.S. 3,867,384 and incorporated herein by reference.

Scheme 7

- 10 One method of preparation of compounds of Formula IIIa is by treatment of a 2-methylthio-4(3H)-quinazolinone of Formula 7 (Z = SMe) with an excess of an amine of Formula HNR^8R^9 at about 150°C to 175°C. A second method is to contact a 2-chloro-4(3H)-quinazolinone of Formula 7 (Z = Cl) with one equivalent of HNR^8R^9 and one equivalent of an acid scavenger, for example triethylamine, or with two
- 15 equivalents of HNR^8R^9 , at a temperature between 60°C and 120°C optionally in the presence of a solvent.


- 20 The preparation of compounds of Formula 7 wherein Z is SMe is described above and in U.S. 3,755,582. The synthesis of compounds of Formula 7 wherein Z is Cl is described in U.S. 3,867,384. Amines of Formula HNR^8R^9 are commercially available or can be prepared by well-known methods (March, J. *Advanced Organic Chemistry*; 3rd ed., John Wiley: New York, (1985), p 1153).

In addition to the methods described above, compounds of Formula Ia and IIa can be prepared by displacement of the 2-chlorine in the appropriate 4(3H)-quinazolinone,

rather than by displacement of the 2-SCH₃ group (Scheme 1) or S-alkylation of the thiocarbonyl (Scheme 4).


As described above for compounds of Formula Ib and IIc, quinazolinethiones of Formula IIIb can be prepared by treatment of the corresponding quinazolinone with P₂S₅ or Lawesson's reagent (Scheme 8).

Scheme 8



Alternatively, 4(3H)-quinazolinones and quinazolinethiones of Formulae IIIc and IIIe, compounds of Formula III wherein R⁸ = -C(=O)R¹², can be prepared by acylation of the corresponding quinazolinones or quinazolinethione wherein R⁸ = H (Formula IIIc) as illustrated in Scheme 9.

Scheme 9



The quinazolinones of Formula IIIc can be treated with an acylating agent of Formula R¹⁰C(=O)L wherein L is an appropriate leaving group such as chlorine or OC(=O)(H or C₁-C₄ alkyl). In a similar fashion, compounds of Formula III wherein R⁸

is $-C(=O)NHR^{12}$ (Formula IIIe) can be prepared by condensing quinazolinones of Formula IIIc with isocyanates of Formula $R^{14}N=C=O$ using well known procedures.

Salts of compounds of Formulae I, II, and III can be formed by treating the free base of the corresponding compound with strong acids such as hydrochloric or sulfuric acid. Salts can also be prepared by alkylation of a tertiary amine group in the molecule to form, for example, the trialkylammonium salt. *N*-Oxides of compounds of Formulae I, II, and III can be made by oxidizing the corresponding reduced nitrogen compound with a strong oxidizing agent such as *meta*-chloroperoxybenzoic acid.

EXAMPLE 1

Synthesis of 6-Bromo-3-propyl-2-propyloxy-4(3H)-quinazolinone

All reactions were conducted under a nitrogen atmosphere.

Step A

To a solution of 200 mL of ethanol containing 37 g of 2-amino-5-bromobenzoic acid was added dropwise 17.72 mL of *n*-propyl isothiocyanate with stirring. The mixture was heated at reflux for 8 h, allowed to cool to room temperature and stirred for approximately 60 h. The mixture was then cooled to approximately 5°C and filtered to obtain 15.42 g of an off-white solid.

Step B

To a solution containing 15.4 g of the product of Step A dissolved in 100 mL of 10% propanolic sodium hydroxide was added 3.2 mL of iodomethane with stirring. The mixture was stirred at room temperature for 10 min, then heated at reflux for 1.5 h, and then allowed to cool to room temperature and stirred overnight. The reaction mixture was filtered to obtain 11.47 g of a white solid. The white solid was purified by column chromatography on silica gel eluting with hexane and then 9:1 hexane:ethyl acetate. Collection and evaporation of those fractions containing the least polar component (according to thin layer chromatography, 6:1 hexane/ethyl acetate mixture as the development solvent) yielded 6.55 g of a white solid, m.p. 97-99°C.

Step C

To 150 mL of propanol cooled to approximately -60°C was added 0.83 g of NaH (60% active in oil) with stirring. To this mixture at -60°C was added 6.5 g of the purified product obtained in Step B. The mixture was allowed to warm to room temperature and stirred for approximately 48 h to yield a clear solution. The reaction solution was poured into water and extracted twice with diethyl ether. The ether extracts were washed twice with water, dried over magnesium sulfate, filtered and the filtrate was then evaporated to yield 10.3 g of an oil. Thin layer chromatography indicated starting material and desired product were both present.

Step D

To propanol cooled to -50°C was added 0.60 g of NaH (60% active in oil) with stirring. To this mixture at -40°C was added the product of Step C and the mixture was allowed to warm to room temperature and stirred for approximately 72 h. The mixture was then heated at reflux for 30 min, cooled to room temperature, poured into water and extracted twice with diethyl ether. The combined ether extracts were washed three times with water, dried over magnesium sulfate, filtered and the filtrate was evaporated to yield an oil. The oil was purified by column chromatography on silica gel eluting with hexane followed by 9:1 hexane/ethyl acetate. Collection and evaporation of the fractions containing only the least polar component (according to thin layer chromatography on silica gel, 9:1 hexane/ethyl acetate mixture as the development solvent) yielded 4.46 g of the title compound as a white solid, m.p. 57-59°C: ¹H NMR (400 MHz, CDCl₃) δ 8.3 (s,1H), 7.7 (m,1H), 7.3 (m,1H), 4.43 (t,2H), 4.05 (t,2H), 1.85 (m,2H), 1.7 (m,2H), 1.06 (t,3H), 0.97 (t,3H).

EXAMPLE 2Synthesis of 6-Bromo-3-*n*-butyl-2-*n*-propylamino-4(3H)-quinazolinoneStep A

To a solution of 200 mL of ethanol containing 15.15 g of 2-amino-5-bromobenzoic acid was added dropwise 9.3 mL of *n*-butyl isothiocyanate with stirring. To this reaction solution was added 9.77 mL of triethylamine. The reaction solution was heated at reflux for 4 h during which time a solid precipitated. The reaction mixture was cooled to 0°C and filtered to obtain 19.89 g of an off-white solid, m.p. 246-248°C.

Step B

To a solution containing 7 g of the product of Step A suspended in 50 mL of chloroform was added 1.97 mL of sulfonyl chloride with stirring. The solution was heated at reflux for 5 h, then cooled to room temperature. The reaction solution was poured into water and extracted twice with methylene chloride. The organic extracts were dried over magnesium sulfate, filtered and the filtrate was then evaporated to a yellow solid. The solid was purified by column chromatography on silica gel eluting with 6:1 hexane/ethyl acetate. Collection and evaporation of the fractions containing only the second-least polar component (according to thin layer chromatography on silica gel, 4:1 hexane/ethyl acetate mixture as the development solvent) yielded 3.2 g of white solid, m.p. 56-58°C.

Step C

To a solution containing 1.02 g of the purified product obtained in Step B dissolved in 25 mL of tetrahydrofuran was added 0.5 mL of *n*-propylamine. The reaction mixture was stirred for approximately 24 h at room temperature. The reaction was then filtered and the filtrate was evaporated to obtain an oil. The oil was dissolved in diethyl

ether and washed twice with water and once with brine. The ether solution was dried over magnesium sulfate, filtered and the filtrate was then evaporated to yield 0.74 g of the title compound as a white solid, m.p. 71-73°C: ¹H NMR (400 MHz, CDCl₃) 0.97-1.04 (m,6H), 1.45 (m,2H), 1.70 (m,4H), 3.50 (m,2H), 4.00 (t,2H), 4.50 (s,1H),
5 7.24 (d,1H), 7.60 (d,1H), 8.20 (s,1H).

EXAMPLE 3

Synthesis of 6-Bromo-3-*n*-propyl-2-*n*-propylthio-4(3H)-quinazolinone

Step A

To a solution of 150 mL isopropanol containing 29.7 g of 2-amino-5-
10 bromobenzoic acid was added dropwise 15.64 mL of *n*-propyl isothiocyanate with stirring. The reaction mixture was then heated at reflux for 15 h. The reaction mixture was cooled to 0°C and filtered to obtain 9.12 g of an off-white solid.

Step B

To a solution containing 0.34 g of the product of Step A suspended in 20 mL of
15 10% propanolic sodium hydroxide was added 0.22 mL of iodopropane with stirring. The reaction mixture was stirred 1.5 h at room temperature. The reaction was poured into water and extracted twice with methylene chloride. The methylene chloride extractions were washed twice with water, dried over magnesium sulfate, filtered, and the filtrate was then evaporated to yield a white solid. The solid was purified by column
20 chromatography on silica gel eluting with 8:1 hexane/ethyl acetate. Collections and evaporation of the fractions containing only the least-polar component (according to thin layer chromatography on silica gel, 6:1 hexane/ethyl acetate mixture as the development solvent) yielded 0.27 g of the title compound as a white solid, m.p. 65-67°C: ¹H NMR (400 MHz, CDCl₃): δ 0.99-1.10 (m,6H), 1.80 (m,4H), 3.25 (t,2H), 4.10 (t,2H), 7.41
25 (d,1H), 7.78 (d,1H), 8.30 (s,1H).

Using the procedures outlined in Schemes 1-9 and Examples 1-3, the compounds of Tables 1-12 hereinafter can be prepared. The compounds referred to in the Tables which follow are illustrated below:

The following abbreviations are used in the Tables which follow. All alkyl groups
30 are the normal isomers unless indicated otherwise. See structures in Index Tables A-C hereinafter for ring system numbering.

t = tertiary

MeO = methoxy

s = secondary

Pr = propyl

n = normal

CN = cyano

i = iso

c = cyclo

Me = methyl

MeS = methylthio

Et = ethyl

Bu = butyl

Ph = phenyl

TABLE 1

Compounds of Formula I wherein: Q = O, R² = n-Pr, R³ = 6-Br, R⁴ = H, and

<u>R¹</u>	<u>R¹</u>	<u>R¹</u>	<u>R¹</u>
n-Pr	n-Bu	n-pentyl	n-hexyl
n-decyl	i-Pr	i-Bu	s-Bu
c-propyl	c-butyl	c-pentyl	2-butenyl
3-butenyl	2-butyryl	3-butyryl	CF ₃
2-Cl-Et	3-Br-Pr	CH ₂ CH=CHCl	CH ₂ C≡CCl
CH ₂ OCH ₃	CH ₂ OCH ₂ CH ₃	CH ₂ SCH ₃	CH ₂ SCH ₂ CH ₃
CH ₂ CH ₂ SCH ₃	CH ₂ CH ₂ S(O)CH ₃	CH ₂ CH ₂ CH ₂ S(O) ₂ CH ₃	(c-pentyl)CH ₂
CH ₂ CH ₂ OCH ₂ CH=CH ₂	CH ₂ CH ₂ OCH ₂ CH≡CH	(c-hexyl)OCH ₂	(c-pentyl)SCH ₂
CH ₂ CH ₂ SCH ₂ CH=CH ₂	CH ₂ CH ₂ SCH ₂ CH≡CH	CH ₂ OCF ₃	CH ₂ OCH ₂ CH ₂ Cl
CH ₂ OCH ₂ CH=CHCl	CH ₂ OCH ₂ C≡CBr	CH ₂ CH=CHCH ₂ OCH ₃	CH ₂ C≡CCH ₂ OCH ₃
CH ₂ CH=CHCH ₂ SCH ₃	CH ₂ C≡CCH ₂ SCH ₃	CH ₂ CH ₂ Si(CH ₃) ₃	CH ₂ CH ₂ N(CH ₃) ₂
CH ₂ CH ₂ CH ₂ NHCH ₃	CH ₂ CH ₂ NO ₂	CH ₂ CH ₂ CH ₂ CN	PhCH ₂
OCH ₂ CH ₂ CH ₃	OCH ₂ CH ₂ CF ₃	SCH ₂ CH ₃	SCCl ₃
SCH ₂ CH ₂ Cl	NHCH ₂ CH ₂ CH ₃	N(CH ₃)CH ₂ CH ₃	Ph
2-pyridinyl	2-furanyl	2-thienyl	2-naphthyl
5-benzofuranyl	3-benzothienyl	3-quinolinyl	(4-F-Ph)CH ₂

Compounds of Formula I wherein: Q = O, R² = n-Pr, R³ = 6-I, R⁴ = H, and

<u>R¹</u>	<u>R¹</u>	<u>R¹</u>	<u>R¹</u>
n-Pr	n-Bu	n-pentyl	n-hexyl
n-decyl	i-Pr	i-Bu	s-Bu
c-propyl	c-butyl	c-pentyl	2-butenyl
3-butenyl	2-butyryl	3-butyryl	CF ₃
2-Cl-Et	3-Br-Pr	CH ₂ CH=CHCl	CH ₂ C≡CCl
CH ₂ OCH ₃	CH ₂ OCH ₂ CH ₃	CH ₂ SCH ₃	CH ₂ SCH ₂ CH ₃
CH ₂ CH ₂ SCH ₃	CH ₂ CH ₂ S(O)CH ₃	CH ₂ CH ₂ CH ₂ S(O) ₂ CH ₃	(c-pentyl)CH ₂
CH ₂ CH ₂ OCH ₂ CH=CH ₂	CH ₂ CH ₂ OCH ₂ C≡CH	(c-hexyl)OCH ₂	(c-pentyl)SCH ₂
CH ₂ CH ₂ SCH ₂ CH=CH ₂	CH ₂ CH ₂ SCH ₂ C≡CH	CH ₂ OCF ₃	CH ₂ OCH ₂ CH ₂ Cl
CH ₂ OCH ₂ CH=CHCl	CH ₂ OCH ₂ C≡CBr	CH ₂ CH=CHCH ₂ OCH ₃	CH ₂ C≡CCH ₂ OCH ₃
CH ₂ CH=CHCH ₂ SCH ₃	CH ₂ C≡CCH ₂ SCH ₃	CH ₂ CH ₂ Si(CH ₃) ₃	CH ₂ CH ₂ N(CH ₃) ₂
CH ₂ CH ₂ CH ₂ NHCH ₃	CH ₂ CH ₂ NO ₂	CH ₂ CH ₂ CH ₂ CN	(4-F-Ph)CH ₂
OCH ₂ CH ₂ CH ₃	OCH ₂ CH ₂ CF ₃	SCH ₂ CH ₃	SCCl ₃
SCH ₂ CH ₂ Cl	NHCH ₂ CH ₂ CH ₃	N(CH ₃)CH ₂ CH ₃	PhCH ₂
2-pyridinyl	2-furanyl	2-thienyl	2-naphthyl
5-benzofuranyl	3-benzothienyl	3-quinolinyl	(2-Me-Ph)CH ₂ CH ₂

Compounds of Formula I wherein: Q = O, R² = n-Pr, R³ = 6-I, R⁴ = 8-I, and

R ¹	R ¹	R ¹	R ¹
n-Pr	n-Bu	n-pentyl	n-hexyl
n-decyl	i-Pr	i-Bu	s-Bu
c-propyl	c-butyl	c-pentyl	2-butenyl
3-butenyl	2-butyryl	3-butyryl	CF ₃
2-Cl-Et	3-Br-Pr	CH ₂ CH=CHCl	CH ₂ C≡CCl
CH ₂ OCH ₃	CH ₂ OCH ₂ CH ₃	CH ₂ SCH ₃	CH ₂ SCH ₂ CH ₃
CH ₂ CH ₂ SCH ₃	CH ₂ CH ₂ S(O)CH ₃	CH ₂ CH ₂ CH ₂ S(O) ₂ CH ₃	(c-pentyl)CH ₂
CH ₂ CH ₂ OCH ₂ CH=CH ₂	CH ₂ CH ₂ OCH ₂ C≡CH	(c-hexyl)OCH ₂	(c-pentyl)SCH ₂
CH ₂ CH ₂ SCH ₂ CH=CH ₂	CH ₂ CH ₂ SCH ₂ C≡CH	CH ₂ OCF ₃	CH ₂ OCH ₂ CH ₂ Cl
CH ₂ OCH ₂ CH=CHCl	CH ₂ OCH ₂ C≡CBr	CH ₂ CH=CHCH ₂ OCH ₃	CH ₂ C≡CCH ₂ OCH ₃
CH ₂ CH=CHCH ₂ SCH ₃	CH ₂ C≡CCH ₂ SCH ₃	CH ₂ CH ₂ Si(CH ₃) ₃	CH ₂ CH ₂ N(CH ₃) ₂
CH ₂ CH ₂ CH ₂ NHCH ₃	CH ₂ CH ₂ NO ₂	CH ₂ CH ₂ CH ₂ CN	PhCH ₂
OCH ₂ CH ₂ CH ₃	OCH ₂ CH ₂ CF ₃	SCH ₂ CH ₃	SCCl ₃
SCH ₂ CH ₂ Cl	NHCH ₂ CH ₂ CH ₃	N(CH ₃)CH ₂ CH ₃	(2-Me-Ph)CH ₂ CH ₂
2-pyridinyl	2-furanyl	2-thienyl	2-naphthyl
5-benzofuranyl	3-benzothieryl	3-quinolinyl	(4-F-Ph)CH ₂

TABLE 2

Compounds of Formula I wherein: Q = O, R¹ = n-Pr, R³ = 6-Br, R⁴ = H, and

R ²	R ²	R ²	R ²
CH ₂ CH ₂ CH ₂ F	t-Bu	i-Pr	n-Bu
i-Bu	s-Bu	n-pentyl	n-hexyl
n-decyl	c-hexyl	allyl	2-butenyl
3-butenyl	5-decenyl	propargyl	2-butyryl
3-butyryl	CF ₃	CH ₂ CF ₃	CH ₂ CH=CHCl
CH ₂ C≡CBr	CH ₂ OCH ₃	CH ₂ OCH ₂ CH ₃	CH ₂ CH ₂ OCH ₃
CH ₂ SCH ₃	CH ₂ CH ₂ SCH ₃	CH ₂ CH ₂ CH ₂ S(O) ₂ CH ₃	(c-pentyl)CH ₂
2-Cl-Et	CH ₂ CH ₂ OCH ₂ C≡CH	CH ₂ CH ₂ SCH ₂ CH=CH ₂	(c-propyl)OCH ₂
(c-hexyl)SCH ₂	CH ₂ CH ₂ OCF ₃	CH ₂ CH ₂ SCH ₂ C≡CH	CH ₂ CH ₂ CH ₂ CN
CH ₂ CH ₂ Si(CH ₃) ₃	-NHPh	CH ₂ CH ₂ OCH ₂ CCl=CH ₂	CH ₂ OCH ₂ CH ₂ Cl
CH ₂ (4-F-Ph)	-N(CH ₃)Ph	CH ₂ CH ₂ CH ₂ N(CH ₃) ₂	CH ₂ CH ₂ CH ₂ Ph
CH ₂ CH ₂ CH ₂ F	CH ₂ Ph	CH ₂ CH ₂ OCH ₂ CH=CH ₂	CH ₂ CH ₂ Ph
CH ₂ CH ₂ CH ₂ NHCH ₃	CH ₂ CH ₂ NO ₂	-N=CHPh	CH ₂ CH ₂ (4-F-Ph)
-N=CHCH ₂ CH ₂ CH ₃	-N=C(CH ₃) ₂	NHCH ₂ CH ₂ CH ₃	N(CH ₃) ₂

Compounds of Formula I wherein: Q = O, R¹ = n-Pr, R³ = 6-I, R⁴ = H, and

R ²	R ²	R ²	R ²
CH ₂ CH ₂ CH ₂ F	t-Bu	i-Pr	n-Bu
i-Bu	s-Bu	n-pentyl	n-hexyl
n-decyl	c-hexyl	allyl	2-butenyl
3-butenyl	5-decenyl	propargyl	2-butyne
3-butyne	CF ₃	CH ₂ CF ₃	CH ₂ CH=CHCl
CH ₂ C≡CBr	CH ₂ OCH ₃	CH ₂ OCH ₂ CH ₃	CH ₂ CH ₂ OCH ₃
CH ₂ SCH ₃	CH ₂ CH ₂ SCH ₃	CH ₂ CH ₂ CH ₂ S(O) ₂ CH ₃	(c-pentyl)CH ₂
2-Cl-Et	CH ₂ CH ₂ OCH ₂ C≡CH	CH ₂ CH ₂ SCH ₂ CH=CH ₂	(c-propyl)OCH ₂
(c-hexyl)SCH ₂	CH ₂ CH ₂ OCF ₃	CH ₂ CH ₂ SCH ₂ C≡CH	CH ₂ CH ₂ CH ₂ CN
CH ₂ CH ₂ Si(CH ₃) ₃	CH ₂ CH ₂ CO ₂ Et	CH ₂ CH ₂ OCH ₂ CCl=CH ₂	CH ₂ OCH ₂ CH ₂ Cl
Ph	4-Me-Ph	CH ₂ CH ₂ CH ₂ N(CH ₃) ₂	2-F-Ph
4-MeO-Ph	CH ₂ Ph	CH ₂ CH ₂ OCH ₂ CH=CH ₂	CH ₂ CH ₂ Ph
CH ₂ CH ₂ CH ₂ NHCH ₃	CH ₂ CH ₂ NO ₂	-N=CHPh	CH ₂ CH ₂ (4-F-Ph)
-N=CHCH ₂ CH ₂ CH ₃	-N=C(CH ₃) ₂	NHCH ₂ CH ₂ CH ₃	N(CH ₃) ₂
2,4-diCl-Ph	2,4,6-triF-Ph	4-CF ₃ -Ph	2-CN-Ph
CH ₂ (4-F-Ph)	-NHPh	-N(CH ₃)Ph	CH ₂ CH ₂ CH ₂ Ph

Compounds of Formula I wherein: Q = O, R¹ = n-Pr, R³ = 6-I, R⁴ = 8-I, and

R ²	R ²	R ²	R ²
CH ₂ CH ₂ CH ₂ F	t-Bu	i-Pr	n-Bu
i-Bu	s-Bu	n-pentyl	n-hexyl
n-decyl	c-hexyl	allyl	2-butenyl
3-butenyl	5-decenyl	propargyl	2-butyne
3-butyne	CF ₃	CH ₂ CF ₃	CH ₂ CH=CHCl
CH ₂ C≡CBr	CH ₂ OCH ₃	CH ₂ OCH ₂ CH ₃	CH ₂ CH ₂ OCH ₃
CH ₂ SCH ₃	CH ₂ CH ₂ SCH ₃	CH ₂ CH ₂ CH ₂ S(O) ₂ CH ₃	(c-pentyl)CH ₂
2-Cl-Et	CH ₂ CH ₂ OCH ₂ C≡CH	CH ₂ CH ₂ SCH ₂ CH=CH ₂	(c-propyl)OCH ₂
(c-hexyl)SCH ₂	CH ₂ CH ₂ OCF ₃	CH ₂ CH ₂ SCH ₂ C≡CH	CH ₂ CH ₂ CH ₂ CN
CH ₂ CH ₂ Si(CH ₃) ₃	CH ₂ CH ₂ CO ₂ Et	CH ₂ CH ₂ OCH ₂ CCl=CH ₂	CH ₂ OCH ₂ CH ₂ Cl
Ph	4-Me-Ph	CH ₂ CH ₂ CH ₂ N(CH ₃) ₂	2-F-Ph
4-MeO-Ph	CH ₂ Ph	CH ₂ CH ₂ OCH ₂ CH=CH ₂	CH ₂ CH ₂ Ph
CH ₂ CH ₂ CH ₂ NHCH ₃	CH ₂ CH ₂ NO ₂	-N=CHPh	CH ₂ CH ₂ (4-F-Ph)
-N=CHCH ₂ CH ₂ CH ₃	-N=C(CH ₃) ₂	NHCH ₂ CH ₂ CH ₃	N(CH ₃) ₂
2,4-diCl-Ph	2,4,6-triF-Ph	4-CF ₃ -Ph	2-CN-Ph
CH ₂ (4-F-Ph)	-NHPh	-N(CH ₃)Ph	CH ₂ CH ₂ CH ₂ Ph

TABLE 3

Compounds of Formula I wherein Q = O and R¹ = R² = n-Pr, and

R ³	R ⁴	R ³	R ⁴	R ³	R ⁴
6-Cl	H	6-Me	H	6-Me ₃ Si	8-Br
6-Br	8-Me	6-Et	8-Br	6-Me ₂ N	H
6-I	8-Br	6-MeO	H	6-EtNH	H
6-Cl	8-Cl	6-MeS	8-MeO	6-Br	8-Me
6-Br	8-Cl	6-SCH ₂ CH=CH ₂	H	6-Br	8-Et
6-I	8-I	6-S(O) ₂ Me	H	6-i-Pr	H
6-C≡CH	H	6-Br	8-CF ₃	6-Br	8-OCF ₃
6-C≡CH	8-Br	6-CH ₂ C≡CH	H	6-CF ₃ O	H
6-c-propyl	H	6-Br	7-Br	6-CH=CH ₂	H
6-CF ₃	H	6-OCH ₂ CH=CH ₂	H	6-Br	7-Me
6-CH ₂ Br	H	6-Br	5-Me	6-Br	5-Br
6-CH=CHBr	H	6-(c-propyl)CH ₂	H	8-Br	H
6-CH ₃ OCH ₂	H	6-I	8-Me	6-Me	8-Br

TABLE 4

Compounds of Formula I wherein Q = S and

R ¹	R ²	R ³	R ⁴	R ¹	R ²	R ³	R ⁴
n-Pr	n-Pr	6-Br	H	n-Pr	n-Pr	6-Br	8-Me
n-Pr	n-Pr	6-I	8-I	n-Pr	n-Pr	6-C≡CH	H
n-Pr	n-Pr	6-I	H	n-Pr	allyl	6-I	H
n-Pr	n-Pr	6-I	8-I	n-Pr	butyl	6-Br	H
3-butenyl	n-Pr	6-Br	H	n-Pr	butyl	6-I	H
n-Pr	allyl	6-Br	H	n-Pr	allyl	6-Br	H
n-Pr	butyl	6-I	H	n-Pr	butyl	6-Br	H
2-Br-Et	n-Pr	6-I	8-I	n-butyl	n-Pr	6-I	8-I
PhCH ₂	n-Pr	6-Br	H	n-butyl	n-Pr	6-Br	H
2-thienyl	allyl	6-Br	H	2-thienyl	allyl	6-I	H
n-Pr	PhCH ₂	6-I	8-I	n-Pr	PhCH ₂	6-I	H
n-Pr	PhCH ₂ CH ₂	6-Br	H	n-Pr	pentyl	6-Br	H

TABLE 5

Compounds of Formula II wherein: Q = O, n is 0, R⁶ = n-Pr, R³ = 6-Br, R⁴ = H, and

R ⁵	R ⁵	R ⁵	R ⁵
n-Pr	n-Bu	n-pentyl	n-octyl

n-decyl	i-Pr	i-Bu	s-Bu
CH ₂ CH ₂ OCH ₃	propargyl	4-pentynyl	2-butenyl
3-butenyl	2-butylnyl	3-butylnyl	CF ₃
2-Cl-Et	3-Br-Pr	CH ₂ CH=CHCl	CH ₂ C≡CCl
CH ₂ OCH ₃	CH ₂ OCH ₂ CH ₃	CH ₂ SCH ₃	CH ₂ SCH ₂ CH ₃
CH ₂ CH ₂ SCH ₃	CH ₂ CH ₂ S(O)CH ₃	CH ₂ CH ₂ CH ₂ S(O) ₂ CH ₃	(c-pentyl)CH ₂
CH ₂ CH ₂ OCH ₂ CH=CH ₂	CH ₂ CH ₂ OCH ₂ C≡CH	(c-hexyl)OCH ₂	(c-pentyl)SCH ₂
CH ₂ CH ₂ SCH ₂ CH=CH ₂	CH ₂ CH ₂ SCH ₂ CH≡CH	CH ₂ OCF ₃	CH ₂ OCH ₂ CH ₂ Cl
CH ₂ OCH ₂ CH=CHCl	CH ₂ OCH ₂ C≡CBr	CH ₂ CH=CHCH ₂ OCH ₃	CH ₂ C≡CCH ₂ OCH ₃
CH ₂ CH=CHCH ₂ SCH ₃	CH ₂ C≡CCH ₂ SCH ₃	CH ₂ CH ₂ Si(CH ₃) ₃	CH ₂ CH ₂ N(CH ₃) ₂
CH ₂ CH ₂ CH ₂ NHCH ₃	CH ₂ CH ₂ NO ₂	CH ₂ CH ₂ CH ₂ CN	SCCl ₃
OCH ₂ CH ₂ CH ₃	OCH ₂ CH ₂ CF ₃	SCH ₂ CH ₃	Ph
SCH ₂ CH ₂ Cl	NHCH ₂ CH ₂ CH ₃	N(CH ₃)CH ₂ CH ₃	2-naphthyl
4-MeS-Ph	2-furanyl	2-thienyl	4-F-Ph
5-benzofuranyl	3-benzothieryl	2-F-4-Cl-Ph	3-CF ₃ O-Ph
2-F-4-Me-Ph	3-MeO-Ph	4-Ph-Ph	CH ₂ Ph
			4-PhO-Ph

Compounds of Formula II wherein: Q = O, n is 0, R⁶ = n-Pr, R³ = 6-I, R⁴ = H, and

R ⁵	R ⁵	R ⁵	R ⁵
n-Pr	n-Bu	n-pentyl	n-octyl
n-decyl	i-Pr	i-Bu	s-Bu
CH ₂ CH ₂ OCH ₃	propargyl	4-pentynyl	2-butenyl
3-butenyl	2-butylnyl	3-butylnyl	CF ₃
2-Cl-Et	3-Br-Pr	CH ₂ CH=CHCl	CH ₂ C≡CCl
CH ₂ OCH ₃	CH ₂ OCH ₂ CH ₃	CH ₂ SCH ₃	CH ₂ SCH ₂ CH ₃
CH ₂ CH ₂ SCH ₃	CH ₂ CH ₂ S(O)CH ₃	CH ₂ CH ₂ CH ₂ S(O) ₂ CH ₃	(c-pentyl)CH ₂
CH ₂ CH ₂ OCH ₂ CH=CH ₂	CH ₂ CH ₂ OCH ₂ C≡CH	(c-hexyl)OCH ₂	(c-pentyl)SCH ₂
CH ₂ CH ₂ SCH ₂ CH=CH ₂	CH ₂ CH ₂ SCH ₂ C≡CH	CH ₂ OCF ₃	CH ₂ OCH ₂ CH ₂ Cl
CH ₂ OCH ₂ CH=CHCl	CH ₂ OCH ₂ C≡CBr	CH ₂ CH=CHCH ₂ OCH ₃	CH ₂ C≡CCH ₂ OCH ₃
CH ₂ CH=CHCH ₂ SCH ₃	CH ₂ C≡CCH ₂ SCH ₃	CH ₂ CH ₂ Si(CH ₃) ₃	CH ₂ CH ₂ N(CH ₃) ₂
CH ₂ CH ₂ CH ₂ NHCH ₃	CH ₂ CH ₂ NO ₂	CH ₂ CH ₂ CH ₂ CN	CH ₂ Ph
OCH ₂ CH ₂ CH ₃	OCH ₂ CH ₂ CF ₃	SCH ₂ CH ₃	SCCl ₃
SCH ₂ CH ₂ Cl	NHCH ₂ CH ₂ CH ₃	N(CH ₃)CH ₂ CH ₃	Ph
4-MeS-Ph	2-furanyl	2-thienyl	2-naphthyl
5-benzofuranyl	3-benzothieryl	4-Ph-Ph	4-F-Ph
2-F-4-Me-Ph	3-MeO-Ph	2-F-4-Cl-Ph	3-CF ₃ O-Ph
			4-PhO-Ph

TABLE 6

Compounds of Formula II wherein: Q = O, n = 0, R⁵ = n-Pr, R³ = 6-Br, R⁴ = H, and

R ⁶	R ⁶	R ⁶	R ⁶
CH ₂ C(CH ₃)=CH ₂	t-Bu	i-Pr	n-Bu
i-Bu	s-Bu	n-pentyl	n-hexyl
n-decyl	CH ₂ CH(CH ₃)CH ₂ CH ₃	allyl	2-butenyl
3-butenyl	5-heptenyl	propargyl	2-butyne
3-butyne	CF ₃	CH ₂ CF ₃	CH ₂ CH=CHCl
CH ₂ C≡CBr	CH ₂ CH ₂ O(CH ₂) ₂ CH ₃	CH ₂ OCH ₂ CH ₃	CH ₂ CH ₂ OCH ₃
CH ₂ SCH ₃	CH ₂ CH ₂ SCH ₃	CH ₂ CH ₂ CH ₂ S(O) ₂ CH ₃	(c-pentyl)CH ₂
2-Cl-Et	CH ₂ CH ₂ OCH ₂ C≡CH	CH ₂ CH ₂ SCH ₂ CH=CH ₂	(c-propyl)OCH ₂
(c-hexyl)SCH ₂	CH ₂ CH ₂ OCF ₃	CH ₂ CH ₂ SCH ₂ C≡CH	(CH ₂) ₄ CN
CH ₂ CH ₂ Si(CH ₃) ₃	CH ₂ CH ₂ CH ₂ CO ₂ Et	CH ₂ CH ₂ OCH ₂ CCl=CH ₂	CH ₂ OCH ₂ CH ₂ Cl
Ph	4-Me-Ph	CH ₂ CH ₂ CH ₂ N(CH ₃) ₂	2-F-Ph
4-MeO-Ph	(CH ₂) ₄ Ph	CH ₂ CH ₂ OCH ₂ CH=CH ₂	CH ₂ CH ₂ CH ₂ Ph
CH ₂ CH ₂ CH ₂ NHCH ₃	CH ₂ CH ₂ NO ₂	-N=CHPh	CH ₂ CH ₂ (4-F-Ph)
4-Cl-Ph	2-Me-Ph	NHCH ₂ CH ₂ CH ₃	N(CH ₃) ₂
2,4-diCl-Ph	2,4,6-triF-Ph	4-CF ₃ -Ph	2-CN-Ph
	-NHPH	-N(CH ₃)Ph	CH ₂ CH ₂ CH ₂ (4-F-Ph)

Compounds of Formula II wherein: Q = O, n = 0, R⁵ = n-Pr, R³ = 6-I, R⁴ = H, and

R ⁶	R ⁶	R ⁶	R ⁶
CH ₂ C(CH ₃)=CH ₂	t-Bu	i-Pr	n-Bu
i-Bu	s-Bu	n-pentyl	n-hexyl
n-decyl	CH ₂ CH(CH ₃)CH ₂ CH ₃	allyl	2-butenyl
3-butenyl	5-heptenyl	propargyl	2-butyne
3-butyne	CF ₃	CH ₂ CF ₃	CH ₂ CH=CHCl
CH ₂ C≡CBr	CH ₂ CH ₂ O(CH ₂) ₂ CH ₃	CH ₂ OCH ₂ CH ₃	CH ₂ CH ₂ OCH ₃
CH ₂ SCH ₃	CH ₂ CH ₂ SCH ₃	CH ₂ CH ₂ CH ₂ S(O) ₂ CH ₃	(c-pentyl)CH ₂
2-Cl-Et	CH ₂ CH ₂ OCH ₂ C≡CH	CH ₂ CH ₂ SCH ₂ CH=CH ₂	(c-propyl)OCH ₂
(c-hexyl)SCH ₂	CH ₂ CH ₂ OCF ₃	CH ₂ CH ₂ SCH ₂ C≡CH	(CH ₂) ₄ CN
CH ₂ CH ₂ Si(CH ₃) ₃	CH ₂ CH ₂ CH ₂ CO ₂ Et	CH ₂ CH ₂ OCH ₂ CCl=CH ₂	CH ₂ OCH ₂ CH ₂ Cl
Ph	4-Me-Ph	CH ₂ CH ₂ CH ₂ N(CH ₃) ₂	2-F-Ph
4-MeO-Ph	CH ₂ Ph	CH ₂ CH ₂ OCH ₂ CH=CH ₂	CH ₂ CH ₂ CH ₂ Ph
CH ₂ CH ₂ CH ₂ NHCH ₃	CH ₂ CH ₂ NO ₂	-N=CHPh	CH ₂ CH ₂ CH ₂ (4-F-Ph)
4-Cl-Ph	2-Me-Ph	NHCH ₂ CH ₂ CH ₃	N(CH ₃) ₂
2,4-diCl-Ph	2,4,6-triF-Ph	4-CF ₃ -Ph	2-CN-Ph
	-NHPH	-N(CH ₃)Ph	

TABLE 7

Compounds of Formula II wherein Q = O, n = 0, R⁵ = R⁶ = n-Pr, and

R ³	R ⁴	R ³	R ⁴	R ³	R ⁴
6-Cl	H	6-Me	H	6-Me ₃ Si	8-Br
6-Br	8-Me	6-Et	8-Br	6-Me ₂ N	H
6-I	8-Br	6-MeO	H	6-EtNH	H
6-Cl	8-Cl	6-MeS	8-MeO	6-Br	8-Me
6-Br	8-Cl	6-SCH ₂ CH=CH ₂	H	6-Br	8-Et
6-I	8-I	6-S(O) ₂ Me	H	6-i-Pr	H
6-C≡CH	H	6-Br	8-CF ₃	6-Br	8-OCF ₃
6-C≡CH	8-Br	6-CH ₂ C≡CH	H	6-CF ₃ O	H
6-c-propyl	H	6-Br	7-Br	6-CH=CH ₂	H
6-CF ₃	H	6-OCH ₂ CH=CH ₂	H	6-Br	7-Me
6-CH ₂ Br	H	6-Br	5-Me	6-Br	5-Br
6-CH=CHBr	H	6-(c-propyl)CH ₂	H	8-Br	H
6-MeOCH ₂	H	6-I	8-Me	6-Me	8-Br

TABLE 8

Compounds of Formula II wherein Q = O, n = 1

R ⁵	R ⁶	R ³	R ⁴
n-Pr	n-Pr	6-Br	H
n-Pr	n-Pr	6-I	8-I
n-Pr	n-Pr	6-I	H
n-Pr	n-Pr	6-I	8-I
3-butenyl	n-Pr	6-Br	H
n-Pr	allyl	6-Br	H
n-Pr	butyl	6-I	H
2-Br-Et	n-Pr	6-I	8-I
Ph	n-Pr	6-Br	H
4-F-Ph	n-Pr	6-I	H
2-thienyl	butyl	6-Br	H
n-Pr	PhCH ₂ CH ₂ CH ₂	6-Br	H

Compounds of Formula II wherein Q = O, n = 2

R ⁵	R ⁶	R ³	R ⁴
n-Pr	n-Pr	6-Br	H
n-Pr	n-Pr	6-I	8-I
n-Pr	n-Pr	6-I	H
n-Pr	n-Pr	6-I	8-I
3-butenyl	n-Pr	6-Br	H
n-Pr	allyl	6-Br	H
n-Pr	butyl	6-I	H
2-Br-Et	n-Pr	6-I	8-I
Ph	n-Pr	6-Br	H
4-F-Ph	n-Pr	6-I	H
2-thienyl	butyl	6-Br	H
n-Pr	pentyl	6-Br	H

TABLE 9

Compounds of Formula III wherein: Q = O, R⁸ = H, R⁹ = n-Pr, R³ = 6-Br, R⁴ = H, and

R ⁷	R ⁷	R ⁷	R ⁷
n-Pr	n-Bu	n-pentyl	n-hexyl
c-hexyl	i-Pr	i-Bu	s-Bu

c-propyl	c-butyl	c-pentyl	2-butenyl
3-butenyl	2-propynyl	3-pentynyl	CH ₂ CF ₃
2-Cl-Et	3-Br-Pr	CH ₂ CH=CHCl	CH ₂ C≡CCl
CH ₂ OCH ₃	CH ₂ OCH ₂ CH ₃	CH ₂ SCH ₃	CH ₂ SCH ₂ CH ₃
CH ₂ CH ₂ SCH ₃	CH ₂ CH ₂ S(O)CH ₃	CH ₂ CH ₂ CH ₂ S(O) ₂ CH ₃	n-decyl
CH ₂ CH ₂ OCH ₂ CH=CH ₂	CH ₂ CH ₂ OCH ₂ C≡CH	(c-hexyl)OCH ₂	(c-pentyl)SCH ₂
CH ₂ CH ₂ SCH ₂ CH=CH ₂	CH ₂ CH ₂ SCH ₂ C≡CH	CH ₂ OCF ₃	CH ₂ OCH ₂ CH ₂ Cl
CH ₂ OCH ₂ CH=CHCl	CH ₂ OCH ₂ C≡CBr	CH ₂ CH=CHCH ₂ OCH ₃	CH ₂ C≡CCH ₂ OCH ₃
CH ₂ CH=CHCH ₂ SCH ₃	CH ₂ C≡CCH ₂ SCH ₃	CH ₂ CH ₂ Si(CH ₃) ₃	CH ₂ CH ₂ N(CH ₃) ₂
CH ₂ CH ₂ CH ₂ NHCH ₃	CH ₂ CH ₂ NO ₂	CH ₂ CH ₂ CH ₂ CN	SCCl ₃
OCH ₂ CH ₂ CH ₃	OCH ₂ CH ₂ CF ₃	SCH ₂ CH ₃	Ph
SCH ₂ CH ₂ Cl	NHCH ₂ CH ₂ CH ₃	N(CH ₃)CH ₂ CH ₃	2-naphthyl
2-pyridinyl	2-furanyl	2-thienyl	4-F-Ph
5-benzofuranyl	3-benzothienyl	3-quinolinyl	3-CF ₃ O-Ph
2-F-4-Me-Ph	3-MeO-Ph	2-F-4-Cl-Ph	4-Cl-Ph
4-MeS-Ph	4-PhO-Ph	4-Ph-Ph	

Compounds of Formula III wherein: Q = O, R⁸ = H, R⁹ = n-Pr, R³ = 6-I, R⁴ = H, and R⁷

R ⁷	R ⁷	R ⁷	R ⁷
n-Pr	n-Bu	n-pentyl	n-hexyl
c-hexyl	i-Pr	i-Bu	s-Bu
c-propyl	c-butyl	c-pentyl	2-butenyl
3-butenyl	2-propynyl	3-pentynyl	CH ₂ CF ₃
2-Cl-Et	3-Br-Pr	CH ₂ CH=CHCl	CH ₂ C≡CCl
CH ₂ OCH ₃	CH ₂ OCH ₂ CH ₃	CH ₂ SCH ₃	CH ₂ SCH ₂ CH ₃
CH ₂ CH ₂ SCH ₃	CH ₂ CH ₂ S(O)CH ₃	CH ₂ CH ₂ CH ₂ S(O) ₂ CH ₃	n-decyl
CH ₂ CH ₂ OCH ₂ CH=CH ₂	CH ₂ CH ₂ OCH ₂ C≡CH	(c-hexyl)CH ₂	(c-propyl)CH ₂
CH ₂ CH ₂ SCH ₂ CH=CH ₂	CH ₂ CH ₂ SCH ₂ C≡CH	CH ₂ OCF ₃	CH ₂ OCH ₂ CH ₂ Cl
CH ₂ OCH ₂ CH=CHCl	CH ₂ OCH ₂ C≡CBr	CH ₂ CH=CHCH ₂ OCH ₃	CH ₂ C≡CCH ₂ OCH ₃
CH ₂ CH=CHCH ₂ SCH ₃	CH ₂ C≡CCH ₂ SCH ₃	CH ₂ CH ₂ Si(CH ₃) ₃	CH ₂ CH ₂ N(CH ₃) ₂
CH ₂ CH ₂ CH ₂ NHCH ₃	CH ₂ CH ₂ NO ₂	CH ₂ CH ₂ CH ₂ CN	SCCl ₃
OCH ₂ CH ₂ CH ₃	OCH ₂ CH ₂ CF ₃	SCH ₂ CH ₃	Ph
SCH ₂ CH ₂ Cl	NHCH ₂ CH ₂ CH ₃	N(CH ₃)CH ₂ CH ₃	2-naphthyl
2-pyridinyl	2-furanyl	2-thienyl	4-F-Ph
5-benzofuranyl	3-benzothienyl	3-quinolinyl	3-CF ₃ O-Ph
2-F-4-Me-Ph	3-MeO-Ph	2-F-4-Cl-Ph	4-Cl-Ph
4-MeS-Ph	4-PhO-Ph	4-Ph-Ph	

Compounds of Formula III wherein: Q = O, R⁸ = H, R⁹ = n-Pr, R³ = 6-I, R⁴ = 8-I, and

R ⁷	R ⁷	R ⁷	R ⁷
n-Pr	n-Bu	n-pentyl	n-hexyl
c-hexyl	i-Pr	i-Bu	s-Bu
c-propyl	c-butyl	c-pentyl	2-butenyl
3-butenyl	2-propynyl	3-pentynyl	CH ₂ CF ₃
2-Cl-Et	3-Br-Pr	CH ₂ CH=CHCl	CH ₂ C=CCl
CH ₂ OCH ₃	CH ₂ OCH ₂ CH ₃	CH ₂ SCH ₃	CH ₂ SCH ₂ CH ₃
CH ₂ CH ₂ SCH ₃	CH ₂ CH ₂ S(O)CH ₃	CH ₂ CH ₂ CH ₂ S(O) ₂ CH ₃	n-decyl
CH ₂ CH ₂ OCH ₂ CH=CH ₂	CH ₂ CH ₂ OCH ₂ C≡CH	(c-hexyl)CH ₂	(c-propyl)CH ₂
CH ₂ CH ₂ SCH ₂ CH=CH ₂	CH ₂ CH ₂ SCH ₂ C≡CH	CH ₂ OCF ₃	CH ₂ OCH ₂ CH ₂ Cl
CH ₂ OCH ₂ CH=CHCl	CH ₂ OCH ₂ C≡CBr	CH ₂ CH=CHCH ₂ OCH ₃	CH ₂ C≡CCH ₂ OCH ₃
CH ₂ CH=CHCH ₂ SCH ₃	CH ₂ C≡CCH ₂ SCH ₃	CH ₂ CH ₂ Si(CH ₃) ₃	CH ₂ CH ₂ N(CH ₃) ₂
CH ₂ CH ₂ CH ₂ NHCH ₃	CH ₂ CH ₂ NO ₂	CH ₂ CH ₂ CH ₂ CN	NHCH ₃
OCH ₂ CH ₂ CH ₃	OCH ₂ CH ₂ CF ₃	SCH ₂ CH ₃	SCCl ₃
SCH ₂ CH ₂ Cl	NHCH ₂ CH ₂ CH ₃	N(CH ₃)CH ₂ CH ₃	Ph
2-pyridinyl	2-furanyl	2-thienyl	2-naphthyl
5-benzofuranyl	3-benzothieryl	3-quinolinyl	4-F-Ph
2-F-4-Me-Ph	3-MeO-Ph	2-F-4-Cl-Ph	3-CF ₃ O-Ph
4-MeS-Ph	4-PhO-Ph	4-Ph-Ph	4-Cl-Ph

TABLE 10

Compounds of Formula III wherein: Q = O, R⁷ = n-Pr, R⁸ = H, R³ = 6-Br, R⁴ = H, and

R ⁹	R ⁹	R ⁹	R ⁹
Et	t-Bu	i-Pr	n-Bu
i-Bu	s-Bu	n-pentyl	n-hexyl
n-decyl	CH ₂ CH(CH ₃)CH ₂ CH ₃	allyl	2-butenyl
3-butenyl	5-heptenyl	propargyl	2-butyryl
3-butyryl	CH ₂ CH ₂ CH ₂ Cl	CH ₂ CH ₂ CF ₃	CH ₂ CH=CHCl
CH ₂ C≡CBr	(CH ₂) ₂ OCH ₂ CH ₃	CH ₂ OCH ₂ CH ₃	CH ₂ CH ₂ OCH ₃
CH ₂ SCH ₃	CH ₂ CH ₂ SCH ₃	CH ₂ CH ₂ CH ₂ S(O) ₂ CH ₃	(c-pentyl)CH ₂
2-Cl-Et	CH ₂ CH ₂ OCH ₂ C≡CH	CH ₂ CH ₂ SCH ₂ CH=CH ₂	(c-propyl)OCH ₂
(c-hexyl)SCH ₂	CH ₂ CH ₂ OCF ₃	CH ₂ CH ₂ SCH ₂ C≡CH	(CH ₂) ₃ CN
CH ₂ CH ₂ Si(CH ₃) ₃	CH ₂ CH ₂ CO ₂ Et	CH ₂ CH ₂ OCH ₂ CCl=CH ₂	CH ₂ OCH ₂ CH ₂ Cl
-N=CHPh	-NHPh	CH ₂ CH ₂ CH ₂ N(CH ₃) ₂	c-propyl
c-hexyl	-NC(=O)NHPh	CH ₂ CH ₂ OCH ₂ CH=CH ₂	-NC(=S)NHPh
CH ₂ CH ₂ CH ₂ NHCH ₃	CH ₂ CH ₂ NO ₂	-NHCH ₂ CH ₂ CH ₃	N(CH ₃) ₃ ⁺ I ⁻
-N=CHCH ₂ CH ₂ CH ₃	-N=C(CH ₃) ₂	NHCH ₂ CH ₂ CH ₃	N(CH ₃) ₂
-OCH ₂ CH ₂ CH ₃	(CH ₂) ₃ (2,4,6-triF-Ph)	CH ₂ (4-CF ₃ -Ph)	-OCH ₂ CH(CH ₃) ₂

Compounds of Formula III wherein: Q = O, R⁷ = n-Pr, R⁸ = H, R³ = 6-I, R⁴ = H, and

R ⁹	R ⁹	R ⁹	R ⁹
Et	t-Bu	i-Pr	n-Bu
i-Bu	s-Bu	n-pentyl	n-hexyl
n-decyl	CH ₂ CH(CH ₃)CH ₂ CH ₃	allyl	2-butenyl
3-butenyl	5-heptenyl	propargyl	2-butylnyl
3-butylnyl	CH ₂ CH ₂ CH ₂ Cl	CH ₂ CH ₂ CF ₃	CH ₂ CH=CHCl
CH ₂ C≡CBr	(CH ₂) ₂ OCH ₂ CH ₃	CH ₂ OCH ₂ CH ₃	CH ₂ CH ₂ OCH ₃
CH ₂ SCH ₃	CH ₂ CH ₂ SCH ₃	CH ₂ CH ₂ CH ₂ S(O) ₂ CH ₃	(c-pentyl)CH ₂
2-Cl-Et	CH ₂ CH ₂ OCH ₂ C≡CH	CH ₂ CH ₂ SCH ₂ CH=CH ₂	(c-propyl)OCH ₂
(c-hexyl)SCH ₂	CH ₂ CH ₂ OCF ₃	CH ₂ CH ₂ SCH ₂ C≡CH	(CH ₂) ₃ CN
CH ₂ CH ₂ Si(CH ₃) ₃	CH ₂ CH ₂ CO ₂ Et	CH ₂ CH ₂ OCH ₂ CCl=CH ₂	CH ₂ OCH ₂ CH ₂ Cl
-N=CHPh	-NHPh	CH ₂ CH ₂ CH ₂ N(CH ₃) ₂	c-propyl
c-hexyl	-NC(=O)NHPh	CH ₂ CH ₂ OCH ₂ CH=CH ₂	-NC(=S)NHPh
CH ₂ CH ₂ CH ₂ NHCH ₃	CH ₂ CH ₂ NO ₂	-NHCH ₂ CH ₂ CH ₃	N(CH ₃) ₃ ⁺ I ⁻
-N=CHCH ₂ CH ₂ CH ₃	-N=C(CH ₃) ₂	NHCH ₂ CH ₂ CH ₃	N(CH ₃) ₂
-OCH ₂ CH ₂ CH ₃	(CH ₂) ₃ (2,4,6-triF-Ph)	CH ₂ (4-CF ₃ -Ph)	-OCH ₂ CH(CH ₃) ₂

Compounds of Formula III wherein: Q = O, R⁷ = n-Pr, R⁸ = H, R³ = 6-I, R⁴ = 8-I, and

R ⁹	R ⁹	R ⁹	R ⁹
Et	t-Bu	i-Pr	n-Bu
i-Bu	s-Bu	n-pentyl	n-hexyl
n-decyl	CH ₂ CH(CH ₃)CH ₂ CH ₃	allyl	2-butenyl
3-butenyl	5-heptenyl	propargyl	2-butylnyl
3-butylnyl	CH ₂ CH ₂ CH ₂ Cl	CH ₂ CH ₂ CF ₃	CH ₂ CH=CHCl
CH ₂ C≡CBr	(CH ₂) ₂ OCH ₂ CH ₃	CH ₂ OCH ₂ CH ₃	CH ₂ CH ₂ OCH ₃
CH ₂ SCH ₃	CH ₂ CH ₂ SCH ₃	CH ₂ CH ₂ CH ₂ S(O) ₂ CH ₃	(c-pentyl)CH ₂
2-Cl-Et	CH ₂ CH ₂ OCH ₂ C≡CH	CH ₂ CH ₂ SCH ₂ CH=CH ₂	(c-propyl)OCH ₂
(c-hexyl)SCH ₂	CH ₂ CH ₂ OCF ₃	CH ₂ CH ₂ SCH ₂ C≡CH	(CH ₂) ₃ CN
CH ₂ CH ₂ Si(CH ₃) ₃	CH ₂ CH ₂ CO ₂ Et	CH ₂ CH ₂ OCH ₂ CCl=CH ₂	CH ₂ OCH ₂ CH ₂ Cl
-N=CHPh	-NHPh	CH ₂ CH ₂ CH ₂ N(CH ₃) ₂	c-propyl
c-hexyl	-NC(=O)NHPh	CH ₂ CH ₂ OCH ₂ CH=CH ₂	-NC(=S)NHPh
CH ₂ CH ₂ CH ₂ NHCH ₃	CH ₂ CH ₂ NO ₂	-NHCH ₂ CH ₂ CH ₃	N(CH ₃) ₃ ⁺ I ⁻
-N=CHCH ₂ CH ₂ CH ₃	-N=C(CH ₃) ₂	NHCH ₂ CH ₂ CH ₃	N(CH ₃) ₂
-OCH ₂ CH ₂ CH ₃	(CH ₂) ₃ (2,4,6-triF-Ph)	CH ₂ (4-CF ₃ -Ph)	-OCH ₂ CH(CH ₃) ₂
Ph	4-F-Ph	2-Me-Ph	2,4-diCl-Ph

TABLE 11

Compounds of Formula III wherein Q = O, R⁸ = H, R⁷ = R⁹ = n-Pr, and

R ³	R ⁴	R ³	R ⁴	R ³	R ⁴
6-Cl	H	6-Me	H	6-Me ₃ Si	8-Br
6-Br	8-Me	6-Et	8-Br	6-Me ₂ N	H
6-I	8-Br	6-MeO	H	6-EtNH	H
6-Cl	8-Cl	6-MeS	8-MeO	6-Br	8-Me
6-Br	8-Cl	6-SCH ₂ CH=CH ₂	H	6-Br	8-Et
6-I	8-I	6-S(O) ₂ Me	H	6-i-Pr	H
6-C≡CH	H	6-Br	8-CF ₃	6-Br	8-OCF ₃
6-C≡CH	8-Br	6-CH ₂ C≡CH	H	6-CF ₃ O	H
6-c-propyl	H	6-Br	7-Br	6-CH=CH ₂	H
6-CF ₃	H	6-OCH ₂ CH=CH ₂	H	6-Br	7-Me
6-CH ₂ Br	H	6-Br	5-Me	6-Br	5-Br
6-CH=CHBr	H	6-(c-propyl)CH ₂	H	8-Br	H
6-MeOCH ₂	H	6-I	8-Me	6-Me	8-Br

TABLE 12

Compounds of Formula III wherein Q=S, R⁸=H

R ⁷	R ⁹	R ³	R ⁴
n-Pr	n-Pr	6-Br	H
n-Pr	n-Pr	6-I	8-I
n-Pr	n-Pr	6-I	H
n-Pr	n-Pr	6-I	8-I
3-butenyl	n-Pr	6-Br	H
n-Pr	allyl	6-Br	H
n-Pr	Et	6-I	H
2-Br-Et	n-Pr	6-I	8-I
Ph	n-Pr	6-Br	H
4-F-Ph	n-Pr	6-I	H
2-thienyl	Et	6-Br	H
n-Pr	Ph	6-I	8-I
n-Pr	NH(CH ₂) ₂ CH ₃	6-Br	H

Compounds of Formula III wherein Q=O, R⁸=Me

R ⁷	R ⁹	R ³	R ⁴
n-Pr	n-Pr	6-Br	H
n-Pr	n-Pr	6-I	8-I
n-Pr	n-Pr	6-I	H
n-Pr	n-Pr	6-I	8-I
3-butenyl	n-Pr	6-Br	H
n-Pr	allyl	6-Br	H
n-Pr	Et	6-I	H
2-Br-Et	n-Pr	6-I	8-I
Ph	n-Pr	6-Br	H
4-F-Ph	n-Pr	6-I	H
2-thienyl	Et	6-Br	H
n-Pr	Ph	6-I	8-I
n-Pr	(CH ₂) ₃ Cl	6-Br	H

Compounds of Formula III wherein Q = O,

R⁸ = C(=O)OCH₃

R ⁷	R ⁹	R ³	R ⁴
n-Pr	n-Pr	6-Br	H

Compounds of Formula III wherein Q = O,

R⁸ = C(=O)CH₃

R ⁷	R ⁹	R ³	R ⁴
n-Pr	n-Pr	6-Br	H

n-Pr	n-Pr	H	8-I	n-Pr	n-Pr	H	8-I
n-Pr	n-Pr	6-I	H	n-Pr	n-Pr	6-I	H
n-Pr	n-Pr	6-I	8-I	n-Pr	n-Pr	6-I	8-I
3-butenyl	n-Pr	6-Br	H	3-butenyl	n-Pr	6-Br	H
n-Pr	allyl	6-Br	H	n-Pr	allyl	6-Br	H
n-Pr	Et	6-I	H	n-Pr	Et	6-I	H
2-Br-Et	n-Pr	6-I	8-I	2-Br-Et	n-Pr	6-I	8-I
Ph	n-Pr	6-Br	H	Ph	n-Pr	6-Br	H
4-F-Ph	n-Pr	6-I	H	4-F-Ph	n-Pr	6-I	H
2-thienyl	Et	6-Br	H	2-thienyl	Et	6-Br	H
n-Pr	Ph	6-I	8-I	n-Pr	Ph	6-I	8-I
n-Pr	(CH ₂) ₃ Cl	6-Br	H	n-Pr	(CH ₂) ₃ Cl	6-Br	H

Formulation/Utility

Compounds of this invention will generally be used in formulation with an agriculturally suitable composition. The fungicidal compositions of the present invention comprise an effective amount of at least one compound of Formula I, II, or III as defined above and at least one of (a) a surfactant, (b) an organic solvent, and (c) at least one solid or liquid diluent. Useful formulations can be prepared in conventional ways. They include dusts, granules, pellets, solutions, suspensions, emulsions, wettable powders, emulsifiable concentrates, dry flowables and the like. Sprayable formulations can be extended in suitable media and used at spray volumes from about one to several hundred liters per hectare. High strength compositions are primarily used as intermediates for further formulation. The formulations will typically contain effective amounts of active ingredient, diluent and surfactant within the following approximate ranges which add up to 100 weight percent.

	Weight Percent		
	<u>Active Ingredient</u>	<u>Diluent</u>	<u>Surfactant</u>
Wettable Powders	5-90	0-74	1-10
Oil Suspensions, Emulsions, Solutions, (including Emulsifiable Concentrates)	5-50	40-95	0-15
Dusts	1-25	70-99	0-5
Granules, Baits and Pellets	0.01-99	5-99.99	0-15
High Strength Compositions	90-99	0-10	0-2

Typical solid diluents are described in Watkins, et al., *Handbook of Insecticide Dust Diluents and Carriers*, 2nd Ed., Dorland Books, Caldwell, New Jersey. Typical liquid diluents and solvents are described in Marsden, *Solvents Guide*, 2nd Ed., Interscience, New York, (1950). *McCutcheon's Detergents and Emulsifiers Annual*,
5 Allured Publ. Corp., Ridgewood, New Jersey, as well as Sisely and Wood, *Encyclopedia of Surface Active Agents*, Chemical Publ. Co., Inc., New York, (1964), list surfactants and recommended uses. All formulations can contain minor amounts of additives to reduce foam, caking, corrosion, microbiological growth, and the like.

Methods for formulating such compositions are well known. Solutions are
10 prepared by simply mixing the ingredients. Fine solid compositions are made by blending and, usually, grinding as in a hammer mill or fluid energy mill. Water-dispersible granules can be produced by agglomerating a fine powder composition; see for example, Cross et al., *Pesticide Formulations*, Washington, D.C., (1988), pp 251-259. Suspensions are prepared by wet-milling; see, for example, U.S. 3,060,084. Granules
15 and pellets can be made by spraying the active material upon preformed granular carriers or by agglomeration techniques. See Browning, "Agglomeration", *Chemical Engineering*, December 4, 1967, pp 147-148, *Perry's Chemical Engineer's Handbook*, 4th Ed., McGraw-Hill, New York, (1963), pp 8-57 and following, and WO 91/13546. Pellets can be prepared as described in U.S. 4,172,714. Water-dispersible and
20 water-soluble granules can be prepared as taught in DE 3,246,493.

For further information regarding the art of formulation, see U.S. 3,235,361, Col. 6, line 16 through Col. 7, line 19 and Examples 10 through 41; U.S. 3,309,192, Col. 5, line 43 through Col. 7, line 62 and Examples 8, 12, 15, 39, 41, 52, 53, 58, 132, 138-140, 162-164, 166, 167 and 169-182; U.S. 2,891,855, Col. 3, line 66 through Col. 5, line 17
25 and Examples 1-4; Klingman, *Weed Control as a Science*, John Wiley and Sons, Inc., New York, (1961), pp 81-96; and Hance et al., *Weed Control Handbook*, 8th Ed., Blackwell Scientific Publications, Oxford, (1989).

In the following Examples, all percentages are by weight and all formulations are prepared in conventional ways. Compound 1 refers to the compound described in Index
30 Table A hereinafter.

Example A

Wettable Powder

	Compound 1	65.0%
	dodecylphenol polyethylene glycol ether	2.0%
35	sodium ligninsulfonate	4.0%
	sodium silicoaluminate	6.0%
	montmorillonite (calcined)	23.0%.

Example BGranule

	Compound 37	10.0%
5	attapulgate granules (low volatile matter, 0.71/0.30 mm; U.S.S. No. 25-50 sieves)	90.0%.

Example CExtruded Pellet

	Compound 25	25.0%
10	anhydrous sodium sulfate	10.0%
	crude calcium ligninsulfonate	5.0%
	sodium alkyl naphthalenesulfonate	1.0%
	calcium/magnesium bentonite	59.0%.

Example D15 Emulsifiable Concentrate

	Compound 37	20.0%
	blend of oil soluble sulfonates and polyoxyethylene ethers	10.0%
	isophorone	70.0%.

20 The compounds of this invention are useful as plant disease control agents, especially for the control of cereal powdery mildews (e.g., *Erysiphe graminis f. sp. tritici*, the causal agent of wheat powdery mildew). The present invention therefore further comprises a method for controlling plant diseases caused by fungal plant pathogens comprising applying to the plant or portion thereof to be protected, or to the

25 plant seed or seedling to be protected, an effective amount of a compound of Formula I, II, or III or a fungicidal composition containing said compound. The compounds and compositions of this invention provide control of diseases caused by a broad spectrum of fungal plant pathogens in the Basidiomycete, Ascomycete, Oomycete and

30 Deuteromycete classes. They are effective in controlling a broad spectrum of plant diseases, particularly foliar pathogens of ornamental, vegetable, field, cereal, and fruit crops. These pathogens include *Plasmopara viticola*, *Phytophthora infestans*, *Peronospora tabacina*, *Pseudoperonospora cubensis*, *Pythium aphanidermatum*, *Alternaria brassicae*, *Septoria nodorum*, *Cercosporidium personatum*, *Cercospora arachidicola*, *Pseudocercospora herpotrichoides*, *Cercospora beticola*, *Botrytis cinerea*, *Monilinia fructicola*, *Pyricularia oryzae*, *Podosphaera leucotricha*, *Venturia inaequalis*, *Erysiphe graminis*, *Uncinula necatur*, *Puccinia recondita*, *Puccinia graminis*, *Hemileia vastatrix*, *Puccinia striiformis*, *Puccinia arachidis*, *Rhizoctonia solani*, *Sphaerotheca fuliginea*, *Fusarium oxysporum*, *Verticillium dahliae*, *Pythium*

aphanidermatum, *Phytophthora megasperma* and other genera and species closely related to these pathogens.

Compounds of this invention can also be mixed with one or more other insecticides, fungicides, nematocides, bactericides, acaricides, semiochemicals, repellants, attractants, pheromones, feeding stimulants or other biologically active compounds to form a multi-component pesticide giving an even broader spectrum of agricultural protection. Examples of other agricultural protectants with which compounds of this invention can be formulated are: insecticides such as acephate, avermectin B, azinphosmethyl, bifenthrin, biphenate, buprofezin, carbofuran, chlordimeform, chlorpyrifos, cyfluthrin, deltamethrin, diazinon, diflubenzuron, dimethoate, esfenvalerate, fenpropathrin, fenvalerate, fipronil, flucythrinate, flufenprox, fluvalinate, fonophos, isofenphos, malathion, metaldehyde, metha-midophos, methidathion, methomyl, methoprene, methoxychlor, monocrotophos, oxamyl, parathion-methyl, permethrin, phorate, phosalone, phosmet, phosphamidon, pirimicarb, profenofos, rotenone, sulprofos, terbufos, tetrachlorvinphos, thiodicarb, tralomethrin, trichlorfon and triflumuron; fungicides such as benomyl, blasticidin S, bromuconazole, captan, captan, carbendazim, chloroneb, chlorothalonil, copper oxychloride, copper salts, cymoxanil, cyproconazole, cyrodinil, dichloran, diclobutrazol, diclomezine, difenoconazole, diniconazole, dodine, edifenphos, epoxyconazole fenarimol, fenbuconazole, fenpropidine, fenpropimorph, fluquinconazole, flusilazol, flutolanil, flutriafol, folpet, furalaxyl, hexaconazole, ipconazole, iprobenfos, iprodione, isoprothiolane, kasugamycin, mancozeb, maneb, mepronil, metalaxyl, metconazole, myclobutanil, neo-asozin, oxadixyl, penconazole, pencycuron, phosethyl-Al, probenazole, prochloraz, propiconazole, pyrifenoxy, pyrimethanil, pyroquilon, sulfur, tebuconazole, tetraconazole, thiabendazole, thiophanate-methyl, thiuram, triadimefon, triadimenol, tricyclazole, triticonazole, uniconazole, validamycin and vinclozolin; nematocides such as aldoxycarb, fenamiphos and fosthietan; bactericides such as oxytetracycline, streptomycin and tribasic copper sulfate; acaricides such as amitraz, binapacryl, chlorobenzilate, cyhexatin, dicofol, dienochlor, fenbutatin oxide, hexythiazox, oxythioquinox, propargite and tebufenpyrad; and biological agents such as *Bacillus thuringiensis* and baculovirus.

In certain instances, combinations with other fungicides having a similar spectrum of control but a different mode of action will be particularly advantageous for resistance management. Preferred combinations comprise a compound of Formula I, II, or III, and a fungicide selected from the group flusilazole, cyproconazole, tetraconazole, fenpropimorph, fenpropidine, cymoxanil, benomyl, carbendazim, mancozeb, and maneb.

Plant disease control is ordinarily accomplished by applying an effective amount of a compound of this invention either pre- or post-infection, to the portion of the plant to be protected such as the roots, stems, foliage, fruit, seeds, tubers or bulbs, or to the

media (soil or sand) in which the plants to be protected are growing. The compounds can also be applied to the seed to protect the seed and seedling.

Rates of application for these compounds can be influenced by many factors of the environment and should be determined under actual use conditions. Foliage can
5 normally be protected when treated at a rate of from less than 1 g/ha to 5,000 g/ha of active ingredient. Seed and seedlings can normally be protected when seed is treated at a rate of from 0.1 to 10 g per kilogram of seed.

The following TESTS demonstrate the control efficacy of compounds of this invention on specific pathogens. The pathogen control protection afforded by the
10 compounds is not limited, however, to these species. See Index Tables A, B, and C for compound descriptions.

Test compounds were first dissolved in acetone in an amount equal to 3% of the final volume and then suspended at a concentration of 200 ppm in purified water containing 250 ppm of the surfactant Trem® 014 (polyhydric alcohol esters). The
15 resulting test suspensions were then used in the following tests.

TEST A

The test suspension was sprayed to the point of run-off on wheat seedlings. The following day the seedlings were inoculated with a spore dust of *Erysiphe graminis* f. sp. *tritici*, (the causal agent of wheat powdery mildew) and incubated in a growth
20 chamber at 20°C for 7 days, after which disease ratings were made.

TEST B

The test suspension was sprayed to the point of run-off on wheat seedlings. The following day the seedlings were inoculated with a spore suspension of *Puccinia recondita* (the causal agent of wheat leaf rust) and incubated in a saturated atmosphere at
25 20°C for 24 h, and then moved to a growth chamber at 20°C for 6 days, after which disease ratings were made.

TEST C

The test suspension was sprayed to the point of run-off on tomato seedlings. The following day the seedlings were inoculated with a spore suspension of *Phytophthora infestans* (the causal agent of potato and tomato late blight) and incubated in a saturated
30 atmosphere at 20°C for 24 h, and then moved to a growth chamber at 20°C for 5 days, after which disease ratings were made.

TEST D

The test suspension was sprayed to the point of run-off on grape seedlings. The following day the seedlings were inoculated with a spore suspension of *Plasmopara viticola* (the causal agent of grape downy mildew) and incubated in a saturated
35 atmosphere at 20°C for 24 h, moved to a growth chamber at 20°C for 6 days, and then

incubated in a saturated atmosphere at 20°C for 24 h, after which disease ratings were made.


TEST E

The test suspension was sprayed to the point of run-off on cucumber seedlings.

- 5 The following day the seedlings were inoculated with a spore suspension of *Botrytis cinerea* (the causal agent of gray mold on many crops) and incubated in a saturated atmosphere at 20°C for 48 h, and moved to a growth chamber at 20°C for 5 days, after which disease ratings were made.

- 10 In the Tables below, ^a = ¹H NMR data for oils are listed in Index Table D

Index Table A




1a

- 15 Compounds of Formula 1a:

Cmpd No.	R ¹	R ²	R ³	R ⁴	m.p. ^a (°C)
1	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	6-Br	H	57-59
2	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	7-Cl	H	57-60
3	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	5-Cl	H	69-75
4	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	8-Me	H	47-49
5	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	5-Me	H	oil
6	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	6-Me	H	47-50
7	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	6-OMe	7-OMe	112-114
8	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	7-F	H	oil
9	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	7-NO ₂	H	64-66
10	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	6-OMe	H	49-52
11	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	6-Me	8-Me	81-84
12	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	6-C≡CH	H	105-107
13	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	6-F	H	60-62
14	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	6-Cl	H	64-66
15	CH ₂ CH ₂ CH ₃	CH ₂ CH=CH ₂	6-Cl	H	78-80
16	CH ₂ CH ₂ CH ₃	CH ₂ CH=CH ₂	6-Br	H	73-75
17	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	6-Cl	8-Cl	78-80

18	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	6-Br	8-Br	89-94
22	CH ₂ CH ₂ CH ₃	(CH ₂) ₃ CH ₃	6-Br	H	58-59
23	CH ₂ CH ₂ CH ₃	i-Pr	6-Br	H	45-46
25	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	6-I	H	48-49
26	CH ₂ CH ₂ CH ₃	(CH ₂) ₄ CH ₃	6-Br	H	56-57
27	CH ₂ CH ₂ CH ₃	(CH ₂) ₅ CH ₃	6-Br	H	oil
28	CH ₂ CH ₂ CH ₃	i-Pr	6-Cl	H	48-49
29	(CH ₂) ₃ CH ₃	CH ₂ CH ₂ CH ₃	6-Br	H	56-58
30	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH=CH ₂	6-Cl	H	oil
31	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ C(CH ₃) ₃	6-Br	H	70-72
32	CH ₂ CH ₂ CH ₃	(CH ₂) ₃ CH ₂ SCH ₃	6-Br	H	86-91
33	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH=CH ₂	6-Br	H	oil
34	CH ₂ CH ₂ CH ₃	(CH ₂) ₄ CH ₃	6-Cl	H	oil
35	CH ₂ CH ₂ CH ₃	(CH ₂) ₄ CH ₃	6-I	H	47-49
36	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH=CH ₂	6-I	H	43-46
37	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	6-I	8-I	135-138
38	(CH ₂) ₃ CH ₃	(CH ₂) ₃ CH ₃	6-Br	H	oil
39	CH ₂ CH ₂ CH ₃	(CH ₂) ₂ CH ₂ Ph	6-Br	H	72-74
40	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ OCH ₃	6-Br	H	55-57
41	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ N(CH ₃) ₂	6-Br	H	39-42
42	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ N(CH ₃) ₂ · HCl	6-Br	H	215-230
43	(CH ₂) ₃ N(CH ₃) ₂	CH ₂ CH ₂ CH ₃	6-Br	H	oil
44	(CH ₂) ₃ OCH ₃	CH ₂ CH ₂ CH ₃	6-Br	H	61-64
45	CH ₂ CH(CH ₃) ₂	CH ₂ CH ₂ CH ₃	6-Br	H	50-55
46	(c-propyl)CH ₂	CH ₂ CH ₂ CH ₃	6-Br	H	99-101
47	CH(CH ₃)Et	CH ₂ CH ₂ CH ₃	6-Br	H	oil
48	(CH ₂) ₄ CH ₃	(CH ₂) ₄ CH ₃	6-Br	H	oil
49	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	6-NO ₂	H	68-75
50	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	6-C≡C-SiMe ₃	H	76-78
51	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₂ CH ₃	6-I	H	54-57
52	(CH ₂) ₃ CH ₃	(CH ₂) ₃ CH ₃	6-I	H	50-51
53	(CH ₂) ₃ CH ₃	CH ₂ CH ₂ CH ₃	6-I	H	50-52
54	(CH ₂) ₃ SCH ₃	CH ₂ CH ₂ CH ₃	6-Br	H	69-71
55	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ N ⁺ (CH ₃) ₃ I ⁻	6-Br	H	223-225
56	(CH ₂) ₃ N ⁺ (CH ₃) ₃ I ⁻	CH ₂ CH ₂ CH ₃	6-Br	H	200-204
57	(CH ₂) ₃ N(CH ₃) ₂ HCl	CH ₂ CH ₂ CH ₃	6-Br	H	145-150
58	CH ₂ CHBrCH ₂ Br	CH ₂ CH ₂ CH ₃	6-Br	H	118-121
59	CH ₂ CH ₂ (N-1,4-morpholinyl)	CH ₂ CH ₂ CH ₃	6-Br	H	103-105

Index Table B




IIa

Compounds of Formula IIa:

Cmpd No.	<u>R⁵</u>	<u>R⁶</u>	<u>R³</u>	<u>R⁴</u>	m.p. (°C)
60	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	6-I	H	90-92
61	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	6-Br	H	65-67

5

Index Table C



IIIa

Compounds of Formula IIIa:

Cmpd

No.	<u>R⁷</u>	<u>R⁹</u>	<u>R⁸</u>	<u>R³</u>	<u>R⁴</u>	m.p. ^a (°C)
62	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	H	6-Br	H	107-111
63	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	6-Br	H	oil
64	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	H	6-I	H	109-111
65	CH ₂ CH ₂ CH ₃	(CH ₂) ₃ CH ₃	H	6-Br	H	87-88
66	CH ₂ CH ₂ CH ₃	(CH ₂) ₃ CH ₃	H	6-I	H	84-85
67	CH ₂ CH ₂ CH ₃	CH ₂ CH(CH ₃) ₂	H	6-I	H	117-119
68	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	H	6-I	8-I	122-126
69	(CH ₂) ₃ CH ₃	CH ₂ CH ₂ CH ₃	H	6-Br	H	71-73
70	(CH ₂) ₃ CH ₃	(CH ₂) ₃ CH ₃	H	6-Br	H	oil
71	CH ₂ CH ₂ CH ₃	(CH ₂) ₃ CH ₃	H	6-I	8-I	126-131
72	(CH ₂) ₃ CH ₃	(CH ₂) ₃ CH ₃	H	6-I	H	oil
73	(CH ₂) ₃ CH ₃	CH ₂ CH ₂ CH ₃	H	6-I	H	oil
74	(CH ₂) ₃ CH ₃	CH ₂ CH ₂ CH ₃	H	6-I	8-I	116-118
75	(CH ₂) ₃ CH ₃	(CH ₂) ₃ CH ₃	H	6-I	8-I	115-116
76	CH ₂ CH ₂ CH ₃	CH ₂ CH=CH ₂	H	6-F	H	84-88

77	CH ₂ CH ₂ CH ₃	CH ₂ CH=CH ₂	H	6-Br	H	104-106
79	CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH=CH ₂	H	6-Br	H	72-75
80	CH ₂ CH ₂ CH ₃	Ph	H	6-I	8-I	159-162

Index Table D

<u>Cmpd No.</u>	<u>¹H NMR Data^b</u>
5	7.46 (dd,1H), 7.29 (d,1H), 7.04 (d,1H), 4.42 (t,2H), 4.02 (m,2H), 2.84 (s,3H), 1.85 (m,2H), 1.71 (m,2H), 1.06 (t,3H), 0.98 (t,3H).
8	8.17 (dd,1H), 7.09 (dd,1H), 7.00 (dt,1H), 4.43 (t,2H), 4.05 (m,2H), 1.85 (m,2H), 1.73 (m,2H), 1.07 (t,3H), 0.97 (t,3H).
27	0.93-0.99 (2-t,6H), 1.37 (m,4H), 1.48 (m,2H), 1.75 (m,2H), 1.80 (m,2H), 4.05 (t,2H), 4.46 (t,2H), 7.34 (d,1H), 7.70 (d,1H), 8.30 (s,1H).
30	0.94-0.98 (t,3H), 1.70 (m,2H), 2.59 (m,2H), 4.02 (t,2H), 4.53 (t,2H), 5.19 (dd,2H), 5.90 (m,1H), 7.40 (d,1H), 7.59 (d,1H), 8.12 (s,1H).
33	0.93-0.98 (t,3H), 1.70 (m,2H), 2.60 (q,2H), 4.03 (t,2H), 4.51-4.55 (t,2H), 5.20 (dd,2H), 8.29, 8.30 (m,1H).
34	0.95-0.99 (m,6H), 1.41 (m,4H), 1.70 (m,2H), 1.81 (m,2H), 4.05 (t,2H), 4.44-4.48 (t,2H), 7.40 (d,1H), 7.58 (d,1H), 8.13 (s,1H).
38	0.94-1.03 (2-t,6H), 1.40 (m,2H), 1.48 (m,2H), 1.65 (m,2H), 1.80 (m,2H), 4.10 (t,2H), 4.47 (t,2H), 7.34 (d,1H), 7.70 (d,1H), 8.29 (s,1H).
43	2.22 (s,6H), 7.33 (d,1H), 7.71 (d,1H), 8.30 (s,1H).
47	1.45 (d,3H), 7.30 (d,1H), 7.68 (d,1H), 8.29 (s,1H).
48	7.31 (d,1H), 7.69 (d,1H), 8.30 (s,1H).
63	0.88-0.92 (m,9H), 1.59 (m,4H), 1.75 (m,2H), 3.09-3.13 (t,4H), 4.08 (t,2H), 7.38 (d,1H), 7.70 (d,1H), 8.30 (s,1H).
70	0.99 (m,6H), 1.44 (m,4H), 1.66 (m,2H), 3.53 (q,2H), 4.00 (t,2H), 4.49 (s,1H), 7.25 (d,1H), 7.61 (d,1H), 8.10 (s,1H).
72	8.40 (s,1H), 7.89 (d,1H), 7.10 (d,1H), 4.50 (s,1H), 4.0 (t,2H), 3.53 (q,2H), 1.68 (m,4H), 1.45 (m,4H), 0.96-1.01 (m,6H).
73	8.40 (s,1H), 7.79 (d,1H), 7.10 (d,1H), 4.52 (s,1H), 4.0 (t,2H), 3.49 (q,2H), 1.70 (m,4H), 1.43 (m,2H), 0.96-1.02 (m,6H).

^b Unless indicated otherwise, ¹H NMR spectra were obtained in CDCl₃ on a 400 MHz spectrometer. Data are reported in ppm downfield from tetramethylsilane; s = singlet, d = doublet, t = triplet, m = multiplet, dd = doublet of doublets, dt = doublet of triplets.

Results for Tests A-E are given in Table 13. In the table, a rating of 100 indicates 100% disease control and a rating of 0 indicates no disease control (relative to the controls). "-" = not tested.

5

Table 13

<u>Cmpd No.</u>	<u>Test</u> <u>A¹</u>	<u>Test</u> <u>B</u>	<u>Test</u> <u>C</u>	<u>Test</u> <u>D</u>	<u>Test</u> <u>E</u>
1	100	4	24	58	0
2	75	7	0	18	0
3	72	59	0	92	0
4	41	3	0	26	0
5	54	0	23	50	0
6	45	0	0	0	81
7	7	57	23	50	0
8	14	3	0	26	0
9	96	0	0	39	0
10	40	0	0	17	67
12	100	0	0	91	83
13	95 ²	0	0	0	0
14	100	0	0	41	45
15	99 ³	0	0	41	4
16	100 ³	0	0	41	0
17	99 ³	0	33	0	0
18	100 ³	20	20	0	32
22	100	0	0	41	0
23	97	46	0	0	0
25	100 ³	46	0	8	0
26	-	4 ¹	0	6	0
27	100	7	0	18	0
29	97	46	0	0	0
30	100 ³	3	0	26	0
31	38	3	0	26	0
32	100 ³	3	0	26	0
33	100	3	0	81	0
34	100 ³	3	0	68	0
35	100 ³	0	0	0	0
36	100 ³	3	0	50	0

40

37	100 ³	93	26	13	0
38	100 ³	54	66	99	0
39	99 ³	0	0	16	0
40	100 ³	54	100	16	0
41	100	0	23	41	0
42	-	0	23	0	0
43	100 ³	62	45	62	0
44	100 ³	62	0	0	0
45	100 ³	0	0	0	67
46	100 ³	0	0	17	0
47	50 ¹	0	0	0	0
48	92 ³	61	0	0	0
49	36	16	0	56	0
50	99 ³	0	0	56	0
51	100 ³	4	0	56	63
52	100 ³	57	0	10	36
53	100 ³	4	0	83	36
54	95	43	0	10	0
55	59 ³	81	0	74	0
56	57 ³	92	0	17	0
57	91	12	23	99	37
58	98	-	-	-	-
59	100	56	0	8	0
60	-	0	0	7	0
61	100 ³	15	0	33	65
62	99	83	19	98	28
63	97	0	42	100	28
64	100 ³	76	43	0	0
65	100 ³	23 ²	0	96	44
66	99	0	0	0	0
67	89 ³	7	0	18	0
68	100 ³	0	26	13	0
69	94	79	80	89	0
70	97	63	0	100	0
71	100 ³	57	0	56	63
72	100 ³	4	0	91	63
73	100	57	0	72	63
76	99 ³	16	21	9	0

41

77	99	52	44	100	68
79	97 ²	6	0	39	0
80	100 ³	57	0	56	0


¹ Test was run at 10 ppm unless otherwise indicated.

² Test was run at 40 ppm.

³ Test was run at 2 ppm.

CLAIMS

1. A compound of Formula I, II, or III



wherein:

n is 0, 1 or 2;

Q is independently O or S;

- 10 R¹ is C₃-C₁₀ alkyl; C₃-C₅ cycloalkyl; C₄-C₁₀ alkenyl; C₄-C₁₀ alkynyl; C₁-C₁₀ haloalkyl; C₃-C₁₀ haloalkenyl; C₃-C₁₀ haloalkynyl; C₂-C₁₀ alkoxyalkyl; C₂-C₁₀ alkylthioalkyl; C₂-C₁₀ alkylsulfanylalkyl; C₂-C₁₀ alkylsulfonylalkyl; C₅-C₁₀ cycloalkylalkyl; C₄-C₁₀ alkenyloxyalkyl; C₄-C₁₀ alkynyloxyalkyl; C₄-C₁₀ (cycloalkyl)oxyalkyl; C₄-C₁₀ alkenylthioalkyl; C₄-C₁₀ alkynylthioalkyl; C₆-C₁₀ (cycloalkyl)thioalkyl; C₂-C₁₀ haloalkoxyalkyl; C₄-C₁₀ haloalkenyloxyalkyl; C₄-C₁₀ haloalkynyloxyalkyl; C₄-C₁₀ alkoxyalkenyl; C₄-C₁₀ alkoxyalkynyl; C₄-C₁₀ alkylthioalkenyl; C₄-C₁₀ alkylthioalkynyl; C₄-C₁₀ trialkylsilylalkyl; C₁-C₁₀ alkyl substituted with NR¹¹R¹², nitro, cyano, or phenyl optionally substituted with R¹⁴, R¹⁵, and R¹⁶; C₁-C₁₀ alkoxy; C₁-C₁₀ haloalkoxy; C₁-C₁₀ alkylthio; C₁-C₁₀ haloalkylthio; NR¹¹R¹²; or pyridyl, furanyl, thienyl, naphthyl, benzofuranyl, benzothienyl, or quinolinyll each optionally substituted with R¹⁴, R¹⁵, and R¹⁶;
- 15
- 20
- 25 R² is C₃-C₁₀ alkyl; C₆-C₇ cycloalkyl; C₃-C₁₀ alkenyl; C₃-C₁₀ alkynyl; C₁-C₁₀ haloalkyl; C₃-C₁₀ haloalkenyl; C₃-C₁₀ haloalkynyl; C₂-C₁₀ alkoxyalkyl; C₂-C₁₀ alkylthioalkyl; C₂-C₁₀ alkylsulfanylalkyl; C₂-C₁₀ alkylsulfonylalkyl; C₄-C₁₀ cycloalkylalkyl; C₄-C₁₀ alkenyloxyalkyl; C₄-C₁₀ alkynyloxyalkyl; C₄-C₁₀ (cycloalkyl)oxyalkyl; C₄-C₁₀ alkenylthioalkyl; C₄-C₁₀ alkynylthioalkyl; C₆-C₁₀ (cycloalkyl)thioalkyl; C₂-C₁₀ haloalkoxyalkyl; C₄-C₁₀ haloalkenyloxyalkyl; C₄-C₁₀ haloalkynyloxyalkyl; C₄-C₁₀ alkoxyalkenyl; C₄-C₁₀ alkoxyalkynyl; C₄-C₁₀ alkylthioalkenyl; C₄-C₁₀ alkylthioalkynyl; C₄-C₁₀ trialkylsilylalkyl; C₃-C₁₀ cyanoalkyl; C₂-C₁₀ nitroalkyl; C₁-C₁₀ alkyl substituted with CO₂R¹¹, NR¹¹R¹², or phenyl
- 30

optionally substituted with R^{13} , R^{15} , and R^{16} ; phenyl optionally substituted with R^{13} , R^{15} , and R^{16} ; $-N=CR^{11}R^{11}$; or $-NR^{11}R^{12}$; or

R^1 and R^2 are taken together to form $-\text{CH}_2(\text{CH}_2)_m\text{CH}_2-$;

m is 1-4;

5 R^3 is halogen; $\text{C}_1\text{-C}_8$ alkyl; $\text{C}_3\text{-C}_8$ cycloalkyl; $\text{C}_2\text{-C}_8$ alkenyl; $\text{C}_2\text{-C}_8$ alkynyl; $\text{C}_1\text{-C}_8$ haloalkyl; $\text{C}_3\text{-C}_8$ haloalkenyl; $\text{C}_3\text{-C}_8$ haloalkynyl; $\text{C}_1\text{-C}_8$ alkoxy; $\text{C}_1\text{-C}_8$ haloalkoxy; $\text{C}_3\text{-C}_8$ alkenyloxy; $\text{C}_3\text{-C}_8$ alkynyloxy; $\text{C}_1\text{-C}_8$ alkylthio; $\text{C}_3\text{-C}_8$ alkenylthio; $\text{C}_3\text{-C}_8$ alkynylthio; $\text{C}_1\text{-C}_8$ alkylsulfinyl; $\text{C}_1\text{-C}_8$ alkylsulfonyl; $\text{C}_2\text{-C}_8$ alkoxyalkyl; $\text{C}_2\text{-C}_8$ alkylthioalkyl; $\text{C}_2\text{-C}_8$ alkylsulfinylalkyl; $\text{C}_2\text{-C}_8$ alkylsulfonylalkyl; $\text{C}_4\text{-C}_8$ cycloalkylalkyl; $\text{C}_3\text{-C}_8$ trialkylsilyl; nitro; $\text{NR}^{11}R^{12}$; $\text{C}_5\text{-C}_8$ trialkylsilylalkynyl; or phenyl optionally substituted with at least one R^{13} ;

R^4 is hydrogen; halogen; $\text{C}_1\text{-C}_4$ alkyl; $\text{C}_1\text{-C}_4$ haloalkyl; $\text{C}_1\text{-C}_4$ alkoxy; or $\text{C}_1\text{-C}_4$ haloalkoxy;

15 R^5 is $\text{C}_3\text{-C}_5$ alkyl; $\text{C}_7\text{-C}_{10}$ alkyl; $\text{C}_4\text{-C}_7$ alkenyl; $\text{C}_3\text{-C}_5$ alkynyl; $\text{C}_1\text{-C}_{10}$ haloalkyl; $\text{C}_5\text{-C}_{10}$ haloalkenyl; $\text{C}_3\text{-C}_{10}$ haloalkynyl; $\text{C}_2\text{-C}_{10}$ alkoxyalkyl other than methoxypropyl; $\text{C}_2\text{-C}_{10}$ alkylthioalkyl; $\text{C}_2\text{-C}_{10}$ alkylsulfinylalkyl; $\text{C}_2\text{-C}_{10}$ alkylsulfonylalkyl; $\text{C}_4\text{-C}_{10}$ cycloalkylalkyl; $\text{C}_4\text{-C}_{10}$ alkenyloxyalkyl; $\text{C}_4\text{-C}_{10}$ alkynyloxyalkyl; $\text{C}_4\text{-C}_{10}$ (cycloalkyl)oxyalkyl; $\text{C}_4\text{-C}_{10}$ alkenylthioalkyl; $\text{C}_4\text{-C}_{10}$ alkynylthioalkyl; $\text{C}_6\text{-C}_{10}$ (cycloalkyl)thioalkyl; $\text{C}_2\text{-C}_{10}$ haloalkoxyalkyl; $\text{C}_4\text{-C}_{10}$ haloalkenyloxyalkyl; $\text{C}_4\text{-C}_{10}$ haloalkynyloxyalkyl; $\text{C}_4\text{-C}_{10}$ alkoxyalkenyl; $\text{C}_4\text{-C}_{10}$ alkoxyalkynyl; $\text{C}_4\text{-C}_{10}$ alkylthioalkenyl; $\text{C}_4\text{-C}_{10}$ alkylthioalkynyl; $\text{C}_4\text{-C}_{10}$ trialkylsilylalkyl; $\text{C}_1\text{-C}_{10}$ alkyl substituted with $\text{NR}^{11}R^{12}$, nitro, or phenyl optionally substituted with at least one of R^{14} , R^{15} , and R^{16} ; $\text{C}_2\text{-C}_{10}$ alkyl substituted with cyano; $\text{C}_1\text{-C}_{10}$ alkoxy; $\text{C}_1\text{-C}_{10}$ haloalkoxy; $\text{C}_1\text{-C}_{10}$ alkylthio; $\text{C}_1\text{-C}_{10}$ haloalkylthio; $\text{NR}^{11}R^{12}$; or phenyl, furanyl, thienyl, naphthyl, benzofuranyl, or benzothienyl each optionally substituted with R^{14} , R^{15} , and R^{16} ;

25 R^6 is $\text{C}_3\text{-C}_{10}$ alkyl; $\text{C}_3\text{-C}_7$ alkenyl; $\text{C}_3\text{-C}_{10}$ alkynyl; $\text{C}_1\text{-C}_{10}$ haloalkyl; $\text{C}_3\text{-C}_{10}$ haloalkenyl; $\text{C}_3\text{-C}_{10}$ haloalkynyl; $\text{C}_3\text{-C}_{10}$ alkoxyalkyl other than propoxymethyl; $\text{C}_2\text{-C}_{10}$ alkylthioalkyl; $\text{C}_2\text{-C}_{10}$ alkylsulfinylalkyl; $\text{C}_2\text{-C}_{10}$ alkylsulfonylalkyl; $\text{C}_4\text{-C}_{10}$ cycloalkylalkyl; $\text{C}_4\text{-C}_{10}$ alkenyloxyalkyl; $\text{C}_4\text{-C}_{10}$ alkynyloxyalkyl; $\text{C}_4\text{-C}_{10}$ (cycloalkyl)oxyalkyl; $\text{C}_4\text{-C}_{10}$ alkenylthioalkyl; $\text{C}_4\text{-C}_{10}$ alkynylthioalkyl; $\text{C}_6\text{-C}_{10}$ (cycloalkyl)thioalkyl; $\text{C}_2\text{-C}_{10}$ haloalkoxyalkyl; $\text{C}_4\text{-C}_{10}$ haloalkenyloxyalkyl; $\text{C}_4\text{-C}_{10}$ haloalkynyloxyalkyl; $\text{C}_4\text{-C}_{10}$ alkoxyalkenyl; $\text{C}_4\text{-C}_{10}$ alkoxyalkynyl; $\text{C}_4\text{-C}_{10}$ alkylthioalkenyl; $\text{C}_4\text{-C}_{10}$ alkylthioalkynyl; $\text{C}_4\text{-C}_{10}$ trialkylsilylalkyl; $\text{C}_5\text{-C}_{10}$ cyanoalkyl; $\text{C}_2\text{-C}_{10}$ nitroalkyl; or $\text{C}_3\text{-C}_{10}$ alkyl substituted with CO_2R^{11} , $\text{NR}^{11}R^{12}$, or phenyl

optionally substituted with R^{13} , R^{15} , and R^{16} ; or phenyl optionally substituted with R^{13} , R^{15} , and R^{16} ; or

R^5 and R^6 are taken together to form $-\text{CH}_2(\text{CH}_2)_m\text{CH}_2-$;

R^7 is $\text{C}_3\text{-C}_{10}$ alkyl; $\text{C}_3\text{-C}_7$ cycloalkyl; $\text{C}_4\text{-C}_7$ alkenyl; propynyl; $\text{C}_5\text{-C}_{10}$ alkynyl; $\text{C}_2\text{-C}_{10}$ haloalkyl; $\text{C}_3\text{-C}_{10}$ haloalkenyl; $\text{C}_3\text{-C}_{10}$ haloalkynyl; $\text{C}_2\text{-C}_{10}$ alkoxyalkyl; $\text{C}_2\text{-C}_{10}$ alkylthioalkyl; $\text{C}_2\text{-C}_{10}$ alkylsulfinylalkyl; $\text{C}_2\text{-C}_{10}$ alkylsulfonylalkyl; $\text{C}_4\text{-C}_{10}$ alkenyloxyalkyl; $\text{C}_4\text{-C}_{10}$ alkynyloxyalkyl; $\text{C}_4\text{-C}_{10}$ (cycloalkyl)oxyalkyl; $\text{C}_4\text{-C}_{10}$ alkenylthioalkyl; $\text{C}_4\text{-C}_{10}$ alkynylthioalkyl; $\text{C}_6\text{-C}_{10}$ (cycloalkyl)thioalkyl; $\text{C}_2\text{-C}_{10}$ haloalkoxyalkyl; $\text{C}_4\text{-C}_{10}$ haloalkenyloxyalkyl; $\text{C}_4\text{-C}_{10}$ haloalkynyloxyalkyl; $\text{C}_4\text{-C}_{10}$ alkoxyalkenyl; $\text{C}_4\text{-C}_{10}$ alkoxyalkynyl; $\text{C}_4\text{-C}_{10}$ alkylthioalkenyl; $\text{C}_4\text{-C}_{10}$ alkylthioalkynyl; $\text{C}_4\text{-C}_{10}$ trialkylsilylalkyl; $\text{C}_1\text{-C}_{10}$ alkyl substituted with $\text{NR}^{11}\text{R}^{12}$ or nitro; $\text{C}_2\text{-C}_{10}$ alkyl substituted with cyano; $\text{C}_1\text{-C}_{10}$ alkoxy; $\text{C}_1\text{-C}_{10}$ haloalkoxy; $\text{C}_1\text{-C}_{10}$ alkylthio; $\text{C}_1\text{-C}_{10}$ haloalkylthio; $\text{NR}^{12}\text{R}^{17}$; or phenyl, pyridyl, furanyl, thienyl, naphthyl, benzofuranyl, benzothienyl, or quinolinyll each optionally substituted with R^{14} , R^{15} , and R^{16} ;

R^8 is hydrogen; $\text{C}_1\text{-C}_4$ alkyl; or $-\text{C}(=\text{O})\text{R}^{10}$;

R^9 is hydrogen; $\text{C}_2\text{-C}_{10}$ alkyl; $\text{C}_3\text{-C}_7$ cycloalkyl; $\text{C}_3\text{-C}_{10}$ alkenyl; $\text{C}_3\text{-C}_{10}$ alkynyl; $\text{C}_3\text{-C}_{10}$ haloalkyl; $\text{C}_3\text{-C}_{10}$ haloalkenyl; $\text{C}_3\text{-C}_{10}$ haloalkynyl; $\text{C}_3\text{-C}_{10}$ alkoxyalkyl other than butoxyethyl; $\text{C}_2\text{-C}_{10}$ alkylthioalkyl; $\text{C}_2\text{-C}_{10}$ alkylsulfinylalkyl; $\text{C}_2\text{-C}_{10}$ alkylsulfonylalkyl; $\text{C}_4\text{-C}_{10}$ cycloalkylalkyl; $\text{C}_4\text{-C}_{10}$ alkenyloxyalkyl; $\text{C}_4\text{-C}_{10}$ alkynyloxyalkyl; $\text{C}_4\text{-C}_{10}$ (cycloalkyl)oxyalkyl; $\text{C}_4\text{-C}_{10}$ alkenylthioalkyl; $\text{C}_4\text{-C}_{10}$ alkynylthioalkyl; $\text{C}_6\text{-C}_{10}$ (cycloalkyl)thioalkyl; $\text{C}_2\text{-C}_{10}$ haloalkoxyalkyl; $\text{C}_4\text{-C}_{10}$ haloalkenyloxyalkyl; $\text{C}_4\text{-C}_{10}$ haloalkynyloxyalkyl; $\text{C}_4\text{-C}_{10}$ alkoxyalkenyl; $\text{C}_4\text{-C}_{10}$ alkoxyalkynyl; $\text{C}_4\text{-C}_{10}$ alkylthioalkenyl; $\text{C}_4\text{-C}_{10}$ alkylthioalkynyl; $\text{C}_4\text{-C}_{10}$ trialkylsilylalkyl; $\text{C}_1\text{-C}_{10}$ alkyl substituted with $\text{NR}^{11}\text{R}^{12}$; $\text{C}_4\text{-C}_{10}$ cyanoalkyl; $\text{C}_2\text{-C}_{10}$ nitroalkyl; $\text{C}_1\text{-C}_8$ alkyl substituted with CO_2R^{11} ; pyridyl, furanyl, thienyl, or naphthyl each optionally substituted with R^{14} , R^{15} , and R^{16} ; $-\text{N}=\text{CR}^{11}\text{R}^{11}$; $-\text{NR}^{12}\text{R}^{17}$; $-\text{OR}^{12}$; or $-\text{NC}(=\text{Q})\text{NR}^{11}\text{R}^{12}$; or R^3 and R^4 are both iodine and R^9 is phenyl optionally substituted with R^{14} , R^{15} , and R^{16} ; or

R^7 and R^9 are taken together to form $-\text{CH}_2(\text{CH}_2)_m\text{CH}_2-$;

R^{10} is hydrogen; $\text{C}_1\text{-C}_4$ alkyl; $\text{C}_1\text{-C}_4$ alkoxy; or $\text{NR}^{11}\text{R}^{12}$;

R^{11} is independently hydrogen; $\text{C}_1\text{-C}_4$ alkyl; or phenyl optionally substituted with at least one R^{13} ;

R^{12} is independently hydrogen; $\text{C}_1\text{-C}_8$ alkyl; or phenyl optionally substituted with at least one R^{13} ; or

R¹¹ and R¹² are taken together to form -CH₂CH₂CH₂CH₂-, -CH₂(CH₂)₃CH₂-,
-CH₂CH₂OCH₂CH₂-, -CH₂CH(Me)CH₂CH(Me)CH₂-, or
-CH₂CH(Me)OCH(Me)CH₂-;

R¹³ is independently halogen; C₁-C₄ alkyl; C₁-C₄ alkoxy; C₁-C₄ haloalkyl; nitro;
or cyano;

R¹⁴ is independently C₁-C₆ alkyl; C₁-C₆ alkoxy; C₁-C₆ haloalkyl; halogen; C₂-C₈
alkynyl; C₁-C₆ thioalkyl; phenyl or phenoxy each optionally substituted with
at least one R¹³; cyano; nitro; C₁-C₆ haloalkoxy; C₁-C₆ haloalkylthio; C₂-C₆
alkenyl; C₂-C₆ haloalkenyl; acetyl; CO₂Me; or N(C₁-C₂ alkyl)₂;

R¹⁵ is independently methyl; ethyl; methoxy; methylthio; halogen; or
trifluoromethyl;

R¹⁶ is independently halogen; and

R¹⁷ is independently C₁-C₈ alkyl; or phenyl optionally substituted with at least one
R¹³; or

an *N*-oxide or agriculturally-suitable salt thereof.

2. A compound of Formula I of Claim 1 wherein:

Q is O;

R¹ is C₃-C₈ alkyl; C₄-C₈ alkenyl; C₄-C₈ alkynyl; C₁-C₈ haloalkyl; C₃-C₈
haloalkenyl; C₂-C₈ alkoxyalkyl; C₂-C₈ alkylthioalkyl; C₅-C₈ cycloalkylalkyl;
C₂-C₈ alkyl substituted with cyano; C₁-C₈ alkoxy; C₁-C₈ haloalkoxy; C₁-C₈
alkylthio; or C₄-C₈ alkenyloxyalkyl; or pyridyl, furanyl, or thienyl each
optionally substituted with R¹⁴ and R¹⁵;

R² is C₃-C₈ alkyl; C₃-C₈ alkenyl; C₃-C₈ alkynyl; C₁-C₈ haloalkyl; C₃-C₈
haloalkenyl; C₃-C₈ alkoxyalkyl; C₂-C₈ alkylthioalkyl; C₄-C₈ cycloalkylalkyl;
C₃-C₈ cyanoalkyl; C₄-C₈ alkenyloxyalkyl; or phenyl optionally substituted
with R¹³;

R³ is halogen; C₁-C₈ alkyl; C₂-C₈ alkynyl; C₃-C₈ cycloalkyl; C₁-C₈ haloalkyl;
C₁-C₈ alkoxy; C₁-C₈ haloalkoxy; C₁-C₈ alkylthio; C₁-C₈ alkylsulfonyl;
C₂-C₈ alkoxyalkyl; C₂-C₈ alkylthioalkyl; C₄-C₈ cycloalkylalkyl; or C₅-C₈
trialkylsilylalkynyl; and

R¹⁴ is methyl; ethyl; methoxy; ethoxy; C₁-C₂ haloalkyl; halogen; acetylenyl;
propargyl; methylthio; ethylthio; cyano; nitro; C₁-C₂ haloalkoxy; vinyl; allyl;
acetyl; CO₂Me; or N(C₁-C₂ alkyl)₂.

3. A compound of Formula II of Claim 1 wherein:

Q is O;

n is 0;

- R^3 is halogen; C_1 - C_8 alkyl; C_2 - C_8 alkynyl; C_3 - C_8 cycloalkyl; C_1 - C_8 haloalkyl; C_1 - C_8 alkoxy; C_1 - C_8 haloalkoxy; C_1 - C_8 alkylthio; C_1 - C_8 alkylsulfonyl; C_2 - C_8 alkoxyalkyl; C_2 - C_8 alkylthioalkyl; C_4 - C_8 cycloalkylalkyl; or C_5 - C_8 trialkylsilylalkynyl;
- 5 R^5 is C_3 - C_5 alkyl; C_4 - C_7 alkenyl; C_3 - C_5 alkynyl; C_1 - C_8 haloalkyl; C_5 - C_8 haloalkenyl; C_2 - C_8 alkoxyalkyl other than methoxypropyl; C_2 - C_8 alkylthioalkyl; C_4 - C_8 cycloalkylalkyl; C_2 - C_8 alkyl substituted with cyano; C_1 - C_8 alkoxy; C_1 - C_8 haloalkoxy; C_1 - C_8 alkylthio; or C_4 - C_8 alkenyloxyalkyl; or phenyl, furanyl, or thienyl each optionally substituted with R^{14} and R^{15} ;
- 10 R^6 is C_3 - C_8 alkyl; C_3 - C_7 alkenyl; C_3 - C_8 alkynyl; C_1 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; C_3 - C_8 alkoxyalkyl other than propoxymethyl; C_2 - C_8 alkylthioalkyl; C_4 - C_8 cycloalkylalkyl; C_5 - C_8 cyanoalkyl; C_4 - C_8 alkenyloxyalkyl; phenyl optionally substituted with R^{13} ; or C_3 - C_5 alkyl substituted with phenyl optionally substituted with R^{13} and R^{15} ; and
- 15 R^{14} is methyl; ethyl; methoxy; ethoxy; C_1 - C_2 haloalkyl; halogen; acetylenyl; propargyl; methylthio; ethylthio; cyano; nitro; C_1 - C_2 haloalkoxy; vinyl; allyl; acetyl; CO_2Me ; or $N(C_1-C_2 \text{ alkyl})_2$.

4. A compound of Formula III of Claim 1 wherein:

- 20 Q is O;
- R^3 is halogen; C_1 - C_8 alkyl; C_2 - C_8 alkynyl; C_3 - C_8 cycloalkyl; C_1 - C_8 haloalkyl; C_1 - C_8 alkoxy; C_1 - C_8 haloalkoxy; C_1 - C_8 alkylthio; C_1 - C_8 alkylsulfonyl; C_2 - C_8 alkoxyalkyl; C_2 - C_8 alkylthioalkyl; C_4 - C_8 cycloalkylalkyl; or C_5 - C_8 trialkylsilylalkynyl;
- 25 R^7 is C_3 - C_8 alkyl; C_4 - C_7 alkenyl; propynyl; C_2 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; C_2 - C_8 alkoxyalkyl; C_2 - C_8 alkylthioalkyl; C_2 - C_8 alkyl substituted with cyano; C_1 - C_8 alkoxy; C_1 - C_8 haloalkoxy; C_1 - C_8 alkylthio; or C_4 - C_8 alkenyloxyalkyl; or phenyl, pyridyl, furanyl, or thienyl each optionally substituted with R^{14} and R^{15} ;
- 30 R^9 is C_3 - C_8 alkyl; C_3 - C_8 alkenyl; C_3 - C_8 alkynyl; C_3 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; C_3 - C_8 alkoxyalkyl; C_2 - C_8 alkylthioalkyl; C_4 - C_8 cycloalkylalkyl; C_4 - C_8 cyanoalkyl; C_4 - C_8 alkenyloxyalkyl; $-NR^{12}R^{17}$; or R^3 and R^4 are both iodine and R^9 is phenyl optionally substituted with R^{14} and R^{15} ; and
- 35 R^{14} is methyl; ethyl; methoxy; ethoxy; C_1 - C_2 haloalkyl; halogen; acetylenyl; propargyl; methylthio; ethylthio; cyano; nitro; C_1 - C_2 haloalkoxy; vinyl; allyl; acetyl; CO_2Me ; or $N(C_1-C_2 \text{ alkyl})_2$.

5. A compound of Claim 2 wherein:

R^1 is C_3 - C_8 alkyl; C_4 - C_8 alkenyl; C_4 - C_8 alkynyl; C_3 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; C_3 - C_8 alkoxyalkyl; or thienyl optionally substituted with at least one of R^{14} and R^{15} ;

5 R^2 is C_3 - C_8 alkyl; C_3 - C_8 alkenyl; C_3 - C_8 alkynyl; C_3 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; C_3 - C_8 alkoxyalkyl; or phenyl optionally substituted with R^{13} ;

R^3 is halogen; C_1 - C_4 alkyl; C_1 - C_4 haloalkyl; C_1 - C_4 alkoxy; C_1 - C_4 haloalkoxy; acetylenyl; or trimethylsilylacetylenyl; and

10 R^{14} is methyl; ethyl; methoxy; methylthio; halogen; trifluoromethyl; or $N(C_1$ - C_2 alkyl) $_2$.

6. A compound of Claim 3 wherein:

R^3 is halogen; C_1 - C_4 alkyl; C_1 - C_4 haloalkyl; C_1 - C_4 alkoxy; C_1 - C_4 haloalkoxy; acetylenyl; or trimethylsilylacetylenyl;

15 R^5 is C_3 - C_5 alkyl; C_4 - C_7 alkenyl; C_3 - C_5 alkynyl; C_3 - C_8 haloalkyl; C_5 - C_8 haloalkenyl; C_3 - C_8 alkoxyalkyl; or phenyl or thienyl each optionally substituted with R^{14} and R^{15} ;

R^6 is C_3 - C_8 alkyl; C_3 - C_7 alkenyl; C_3 - C_8 alkynyl; C_3 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; C_3 - C_8 alkoxyalkyl; or phenyl optionally substituted with R^{13} ;

20 and
 R^{14} is methyl; ethyl; methoxy; methylthio; halogen; trifluoromethyl; or $N(C_1$ - C_2 alkyl) $_2$.

7. A compound of Claim 4 wherein:

25 R^3 is halogen; C_1 - C_4 alkyl; C_1 - C_4 haloalkyl; C_1 - C_4 alkoxy; C_1 - C_4 haloalkoxy; acetylenyl; or trimethylsilylacetylenyl;

R^7 is C_3 - C_8 alkyl; C_4 - C_7 alkenyl; propynyl; C_3 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; C_3 - C_8 alkoxy; C_3 - C_8 alkoxyalkyl; or phenyl or thienyl each optionally substituted with R^{14} and R^{15} ;

30 R^9 is C_3 - C_8 alkyl; C_3 - C_8 alkenyl; C_3 - C_8 alkynyl; C_3 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; C_3 - C_8 alkoxyalkyl; $-NR^{12}R^{17}$; or R^3 and R^4 are both iodine and R^9 is phenyl optionally substituted with R^{14} and R^{15} ; and

R^{14} is methyl; ethyl; methoxy; methylthio; halogen; trifluoromethyl; or $N(C_1$ - C_2 alkyl) $_2$.

35

8. A compound of Claim 5 wherein:

R^1 is C_3 - C_8 alkyl; C_4 - C_8 alkenyl; C_4 - C_8 alkynyl; C_3 - C_8 haloalkyl; or C_3 - C_8 haloalkenyl;

R² is C₃-C₈ alkyl; C₃-C₈ alkenyl; C₃-C₈ alkynyl; C₃-C₈ haloalkyl; C₃-C₈ haloalkenyl; or phenyl optionally substituted with R¹³;

R³ is halogen; and

R⁴ is hydrogen or halogen.

5

9. A compound of Claim 6 wherein:

R³ is halogen;

R⁴ is hydrogen or halogen;

R⁵ is C₃-C₅ alkyl; C₄-C₇ alkenyl; C₃-C₅ alkynyl; C₃-C₈ haloalkyl; or C₅-C₈ haloalkenyl; or phenyl optionally substituted with R¹⁴ and R¹⁵; and

10

R⁶ is C₃-C₈ alkyl; C₃-C₇ alkenyl; C₃-C₈ alkynyl; C₃-C₈ haloalkyl; C₃-C₈ haloalkenyl; or phenyl optionally substituted with R¹³.

10. A compound of Claim 7 wherein:

15

R³ is halogen;

R⁴ is hydrogen or halogen;

R⁷ is C₃-C₈ alkyl; C₄-C₇ alkenyl; propynyl; C₃-C₈ haloalkyl; or C₃-C₈ haloalkenyl; or phenyl optionally substituted with R¹⁴ and R¹⁵; and

R⁹ is C₃-C₈ alkyl; C₃-C₈ alkenyl; C₃-C₈ alkynyl; C₃-C₈ haloalkyl; C₃-C₈ haloalkenyl; -NR¹²R¹⁷; or R³ and R⁴ are both iodine and R⁹ is phenyl optionally substituted with R¹⁴ and R¹⁵.

20

11. A compound of Claim 8 which is selected from:

6-bromo-3-propyl-2-propyloxy-4(3H)-quinazolinone;

25

6,8-diiodo-3-propyl-2-propyloxy-4(3H)-quinazolinone; and

6-iodo-3-propyl-2-propyloxy-4(3H)-quinazolinone.




12. A compound of Claim 10 which is

6,8-diiodo-3-propyl-2-(phenylamino)-4(3H)-quinazolinone.

30

13. A method for controlling plant diseases caused by fungal plant pathogens comprising applying to the plant or portion thereof to be protected, to the media in which the plant to be protected is growing, or to the plant seed or seedling to be protected an effective amount of a compound of Formula I, II, or III:

49



wherein:

n is 0, 1 or 2;

5 Q is independently O or S;

R¹ is C₃-C₁₀ alkyl; C₃-C₅ cycloalkyl; C₄-C₁₀ alkenyl; C₄-C₁₀ alkynyl; C₁-C₁₀ haloalkyl; C₃-C₁₀ haloalkenyl; C₃-C₁₀ haloalkynyl; C₂-C₁₀ alkoxyalkyl; C₂-C₁₀ alkylthioalkyl; C₂-C₁₀ alkylsulfinylalkyl; C₂-C₁₀ alkylsulfonylalkyl; C₅-C₁₀ cycloalkylalkyl; C₄-C₁₀ alkenyloxyalkyl; C₄-C₁₀ alkynyloxyalkyl; C₄-C₁₀ (cycloalkyl)oxyalkyl; C₄-C₁₀ alkenylthioalkyl; C₄-C₁₀ alkynylthioalkyl; C₆-C₁₀ (cycloalkyl)thioalkyl; C₂-C₁₀ haloalkoxyalkyl; C₄-C₁₀ haloalkenyloxyalkyl; C₄-C₁₀ haloalkynyloxyalkyl; C₄-C₁₀ alkoxyalkenyl; C₄-C₁₀ alkoxyalkynyl; C₄-C₁₀ alkylthioalkenyl; C₄-C₁₀ alkylthioalkynyl; C₄-C₁₀ trialkylsilylalkyl; C₁-C₁₀ alkyl substituted with NR¹¹R¹², nitro, cyano, or phenyl optionally substituted with R¹⁴, R¹⁵, and R¹⁶; C₁-C₁₀ alkoxy; C₁-C₁₀ haloalkoxy; C₁-C₁₀ alkylthio; C₁-C₁₀ haloalkylthio; NR¹¹R¹²; or pyridyl, furanyl, thienyl, naphthyl, benzofuranyl, benzothienyl, or quinolinyll each optionally substituted with R¹⁴, R¹⁵, and R¹⁶;

20 R² is C₃-C₁₀ alkyl; C₆-C₇ cycloalkyl; C₃-C₁₀ alkenyl; C₃-C₁₀ alkynyl; C₁-C₁₀ haloalkyl; C₃-C₁₀ haloalkenyl; C₃-C₁₀ haloalkynyl; C₂-C₁₀ alkoxyalkyl; C₂-C₁₀ alkylthioalkyl; C₂-C₁₀ alkylsulfinylalkyl; C₂-C₁₀ alkylsulfonylalkyl; C₄-C₁₀ cycloalkylalkyl; C₄-C₁₀ alkenyloxyalkyl; C₄-C₁₀ alkynyloxyalkyl; C₄-C₁₀ (cycloalkyl)oxyalkyl; C₄-C₁₀ alkenylthioalkyl; C₄-C₁₀ alkynylthioalkyl; C₆-C₁₀ (cycloalkyl)thioalkyl; C₂-C₁₀ haloalkoxyalkyl; C₄-C₁₀ haloalkenyloxyalkyl; C₄-C₁₀ haloalkynyloxyalkyl; C₄-C₁₀ alkoxyalkenyl; C₄-C₁₀ alkoxyalkynyl; C₄-C₁₀ alkylthioalkenyl; C₄-C₁₀ alkylthioalkynyl; C₄-C₁₀ trialkylsilylalkyl; C₃-C₁₀ cyanoalkyl; C₂-C₁₀ nitroalkyl; C₁-C₁₀ alkyl substituted with CO₂R¹¹, NR¹¹R¹², or phenyl optionally substituted with R¹³, R¹⁵, and R¹⁶; phenyl optionally substituted with R¹³, R¹⁵, and R¹⁶; -N=CR¹¹R¹¹; or -NR¹¹R¹²; or

R¹ and R² are taken together to form -CH₂(CH₂)_mCH₂-;

m is 1-4;

- 5 R^3 is halogen; C_1 - C_8 alkyl; C_3 - C_8 cycloalkyl; C_2 - C_8 alkenyl; C_2 - C_8 alkynyl; C_1 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; C_3 - C_8 haloalkynyl; C_1 - C_8 alkoxy; C_1 - C_8 haloalkoxy; C_3 - C_8 alkenyloxy; C_3 - C_8 alkynyloxy; C_1 - C_8 alkylthio; C_3 - C_8 alkenylthio; C_3 - C_8 alkynylthio; C_1 - C_8 alkylsulfinyl; C_1 - C_8 alkylsulfonyl; C_2 - C_8 alkoxyalkyl; C_2 - C_8 alkylthioalkyl; C_2 - C_8 alkylsulfinylalkyl; C_2 - C_8 alkylsulfonylalkyl; C_4 - C_8 cycloalkylalkyl; C_3 - C_8 trialkylsilyl; nitro; $NR^{11}R^{12}$; C_5 - C_8 trialkylsilylalkynyl; or phenyl optionally substituted with at least one R^{13} ;
- 10 R^4 is hydrogen; halogen; C_1 - C_4 alkyl; C_1 - C_4 haloalkyl; C_1 - C_4 alkoxy; or C_1 - C_4 haloalkoxy;
- 15 R^5 is C_3 - C_5 alkyl; C_7 - C_{10} alkyl; C_4 - C_7 alkenyl; C_3 - C_5 alkynyl; C_1 - C_{10} haloalkyl; C_5 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_2 - C_{10} alkoxyalkyl other than methoxypropyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_4 - C_{10} cycloalkylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_1 - C_{10} alkyl substituted with $NR^{11}R^{12}$, nitro, or phenyl optionally substituted with at least one of R^{14} , R^{15} , and R^{16} ; C_2 - C_{10} alkyl substituted with cyano; C_1 - C_{10} alkoxy; C_1 - C_{10} haloalkoxy; C_1 - C_{10} alkylthio; C_1 - C_{10} haloalkylthio; $NR^{11}R^{12}$; or phenyl, furanyl, thienyl, naphthyl, benzofuranyl, or benzothienyl each optionally substituted with R^{14} , R^{15} , and R^{16} ;
- 20 R^6 is C_3 - C_{10} alkyl; C_3 - C_7 alkenyl; C_3 - C_{10} alkynyl; C_1 - C_{10} haloalkyl; C_3 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_3 - C_{10} alkoxyalkyl other than propoxymethyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_4 - C_{10} cycloalkylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_5 - C_{10} cyanoalkyl; C_2 - C_{10} nitroalkyl; or C_3 - C_{10} alkyl substituted with CO_2R^{11} , $NR^{11}R^{12}$, or phenyl optionally substituted with R^{13} , R^{15} , and R^{16} ; or phenyl optionally substituted with R^{13} , R^{15} , and R^{16} ; or
- 25 R^5 and R^6 are taken together to form $-CH_2(CH_2)_mCH_2-$;

- R^7 is C_3 - C_{10} alkyl; C_3 - C_7 cycloalkyl; C_4 - C_7 alkenyl; propynyl; C_5 - C_{10} alkynyl; C_2 - C_{10} haloalkyl; C_3 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_2 - C_{10} alkoxyalkyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_1 - C_{10} alkyl substituted with $NR^{11}R^{12}$ or nitro; C_2 - C_{10} alkyl substituted with cyano; C_1 - C_{10} alkoxy; C_1 - C_{10} haloalkoxy; C_1 - C_{10} alkylthio; C_1 - C_{10} haloalkylthio; $NR^{12}R^{17}$; or phenyl, pyridyl, furanyl, thienyl, naphthyl, benzofuranyl, benzothienyl, or quinolinyll each optionally substituted with R^{14} , R^{15} , and R^{16} ;
- R^8 is hydrogen; C_1 - C_4 alkyl; or $-C(=O)R^{10}$;
- R^9 is hydrogen; C_2 - C_{10} alkyl; C_3 - C_7 cycloalkyl; C_3 - C_{10} alkenyl; C_3 - C_{10} alkynyl; C_3 - C_{10} haloalkyl; C_3 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_3 - C_{10} alkoxyalkyl other than butoxyethyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_4 - C_{10} cycloalkylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_1 - C_{10} alkyl substituted with $NR^{11}R^{12}$; C_4 - C_{10} cyanoalkyl; C_2 - C_{10} nitroalkyl; C_1 - C_8 alkyl substituted with CO_2R^{11} ; pyridyl, furanyl, thienyl, or naphthyl each optionally substituted with R^{14} , R^{15} , and R^{16} ; $-N=CR^{11}R^{11}$; $-NR^{12}R^{17}$; $-OR^{12}$; or $-NC(=Q)NR^{11}R^{12}$; or R^3 and R^4 are both iodine and R^9 is phenyl optionally substituted with R^{14} , R^{15} , and R^{16} ; or
- R^7 and R^9 are taken together to form $-CH_2(CH_2)_mCH_2-$;
- R^{10} is hydrogen; C_1 - C_4 alkyl; C_1 - C_4 alkoxy; or $NR^{11}R^{12}$;
- R^{11} is independently hydrogen; C_1 - C_4 alkyl; or phenyl optionally substituted with at least one R^{13} ;
- R^{12} is independently hydrogen; C_1 - C_8 alkyl; or phenyl optionally substituted with at least one R^{13} ; or
- R^{11} and R^{12} are taken together to form $-CH_2CH_2CH_2CH_2-$, $-CH_2(CH_2)_3CH_2-$, $-CH_2CH_2OCH_2CH_2-$, $-CH_2CH(Me)CH_2CH(Me)CH_2-$, or $-CH_2CH(Me)OCH(Me)CH_2-$;

R^{13} is independently halogen; C_1 - C_4 alkyl; C_1 - C_4 alkoxy; C_1 - C_4 haloalkyl; nitro; or cyano;

R^{14} is independently C_1 - C_6 alkyl; C_1 - C_6 alkoxy; C_1 - C_6 haloalkyl; halogen; C_2 - C_8 alkynyl; C_1 - C_6 thioalkyl; phenyl or phenoxy each optionally substituted with at least one R^{13} ; cyano; nitro; C_1 - C_6 haloalkoxy; C_1 - C_6 haloalkylthio; C_2 - C_6 alkenyl; C_2 - C_6 haloalkenyl; acetyl; CO_2Me ; or $N(C_1-C_2 \text{ alkyl})_2$;




R^{15} is independently methyl; ethyl; methoxy; methylthio; halogen; or trifluoromethyl;

R^{16} is independently halogen; and

R^{17} is independently C_1 - C_8 alkyl; or phenyl optionally substituted with at least one R^{13} ; or

an *N*-oxide or agriculturally-suitable salt thereof.

14. A fungicidal composition comprising an effective amount of a compound of Formula I, II, or III:



wherein:

n is 0, 1 or 2;

Q is independently O or S;

R^1 is C_3 - C_{10} alkyl; C_3 - C_5 cycloalkyl; C_4 - C_{10} alkenyl; C_4 - C_{10} alkynyl; C_1 - C_{10} haloalkyl; C_3 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_2 - C_{10} alkoxyalkyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_5 - C_{10} cycloalkylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_1 - C_{10} alkyl substituted with $NR^{11}R^{12}$, nitro, cyano, or phenyl optionally substituted with R^{14} , R^{15} , and R^{16} ; C_1 - C_{10} alkoxy; C_1 - C_{10} haloalkoxy; C_1 - C_{10} alkylthio; C_1 - C_{10} haloalkylthio; $NR^{11}R^{12}$; or pyridyl, furanyl, thienyl, naphthyl, benzofuranyl,

benzothienyl, or quinolinyl each optionally substituted with R^{14} , R^{15} , and R^{16} ;

R^2 is C_3 - C_{10} alkyl; C_6 - C_7 cycloalkyl; C_3 - C_{10} alkenyl; C_3 - C_{10} alkynyl; C_1 - C_{10} haloalkyl; C_3 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_2 - C_{10} alkoxyalkyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_4 - C_{10} cycloalkylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_3 - C_{10} cyanoalkyl; C_2 - C_{10} nitroalkyl; C_1 - C_{10} alkyl substituted with CO_2R^{11} , $NR^{11}R^{12}$, or phenyl optionally substituted with R^{13} , R^{15} , and R^{16} ; phenyl optionally substituted with R^{13} , R^{15} , and R^{16} ; $-N=CR^{11}R^{11}$; or $-NR^{11}R^{12}$; or

R^1 and R^2 are taken together to form $-CH_2(CH_2)_mCH_2-$; m is 1-4;

R^3 is halogen; C_1 - C_8 alkyl; C_3 - C_8 cycloalkyl; C_2 - C_8 alkenyl; C_2 - C_8 alkynyl; C_1 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; C_3 - C_8 haloalkynyl; C_1 - C_8 alkoxy; C_1 - C_8 haloalkoxy; C_3 - C_8 alkenyloxy; C_3 - C_8 alkynyloxy; C_1 - C_8 alkylthio; C_3 - C_8 alkenylthio; C_3 - C_8 alkynylthio; C_1 - C_8 alkylsulfinyl; C_1 - C_8 alkylsulfonyl; C_2 - C_8 alkoxyalkyl; C_2 - C_8 alkylthioalkyl; C_2 - C_8 alkylsulfinylalkyl; C_2 - C_8 alkylsulfonylalkyl; C_4 - C_8 cycloalkylalkyl; C_3 - C_8 trialkylsilyl; nitro; $NR^{11}R^{12}$; C_5 - C_8 trialkylsilylalkynyl; or phenyl optionally substituted with at least one R^{13} ;

R^4 is hydrogen; halogen; C_1 - C_4 alkyl; C_1 - C_4 haloalkyl; C_1 - C_4 alkoxy; or C_1 - C_4 haloalkoxy;

R^5 is C_3 - C_5 alkyl; C_7 - C_{10} alkyl; C_4 - C_7 alkenyl; C_3 - C_5 alkynyl; C_1 - C_{10} haloalkyl; C_5 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_2 - C_{10} alkoxyalkyl other than methoxypropyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_4 - C_{10} cycloalkylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_1 - C_{10} alkyl substituted with $NR^{11}R^{12}$, nitro, or phenyl optionally substituted with at least one of R^{14} , R^{15} , and R^{16} ; C_2 - C_{10} alkyl substituted with cyano; C_1 - C_{10} alkoxy; C_1 - C_{10} haloalkoxy; C_1 - C_{10} alkylthio; C_1 - C_{10} haloalkylthio; $NR^{11}R^{12}$; or

phenyl, furanyl, thienyl, naphthyl, benzofuranyl, or benzothienyl each optionally substituted with R¹⁴, R¹⁵, and R¹⁶;

R⁶ is C₃-C₁₀ alkyl; C₃-C₇ alkenyl; C₃-C₁₀ alkynyl; C₁-C₁₀ haloalkyl; C₃-C₁₀ haloalkenyl; C₃-C₁₀ haloalkynyl; C₃-C₁₀ alkoxyalkyl other than propoxymethyl; C₂-C₁₀ alkylthioalkyl; C₂-C₁₀ alkylsulfinylalkyl; C₂-C₁₀ alkylsulfonylalkyl; C₄-C₁₀ cycloalkylalkyl; C₄-C₁₀ alkenyloxyalkyl; C₄-C₁₀ alkynyloxyalkyl; C₄-C₁₀ (cycloalkyl)oxyalkyl; C₄-C₁₀ alkenylthioalkyl; C₄-C₁₀ alkynylthioalkyl; C₆-C₁₀ (cycloalkyl)thioalkyl; C₂-C₁₀ haloalkoxyalkyl; C₄-C₁₀ haloalkenyloxyalkyl; C₄-C₁₀ haloalkynyloxyalkyl; C₄-C₁₀ alkoxyalkenyl; C₄-C₁₀ alkoxyalkynyl; C₄-C₁₀ alkylthioalkenyl; C₄-C₁₀ alkylthioalkynyl; C₄-C₁₀ trialkylsilylalkyl; C₅-C₁₀ cyanoalkyl; C₂-C₁₀ nitroalkyl; or C₃-C₁₀ alkyl substituted with CO₂R¹¹, NR¹¹R¹², or phenyl optionally substituted with R¹³, R¹⁵, and R¹⁶; or phenyl optionally substituted with R¹³, R¹⁵, and R¹⁶; or

R⁵ and R⁶ are taken together to form -CH₂(CH₂)_mCH₂-;

R⁷ is C₃-C₁₀ alkyl; C₃-C₇ cycloalkyl; C₄-C₇ alkenyl; propynyl; C₅-C₁₀ alkynyl; C₂-C₁₀ haloalkyl; C₃-C₁₀ haloalkenyl; C₃-C₁₀ haloalkynyl; C₂-C₁₀ alkoxyalkyl; C₂-C₁₀ alkylthioalkyl; C₂-C₁₀ alkylsulfinylalkyl; C₂-C₁₀ alkylsulfonylalkyl; C₄-C₁₀ alkenyloxyalkyl; C₄-C₁₀ alkynyloxyalkyl; C₄-C₁₀ (cycloalkyl)oxyalkyl; C₄-C₁₀ alkenylthioalkyl; C₄-C₁₀ alkynylthioalkyl; C₆-C₁₀ (cycloalkyl)thioalkyl; C₂-C₁₀ haloalkoxyalkyl; C₄-C₁₀ haloalkenyloxyalkyl; C₄-C₁₀ haloalkynyloxyalkyl; C₄-C₁₀ alkoxyalkenyl; C₄-C₁₀ alkoxyalkynyl; C₄-C₁₀ alkylthioalkenyl; C₄-C₁₀ alkylthioalkynyl; C₄-C₁₀ trialkylsilylalkyl; C₁-C₁₀ alkyl substituted with NR¹¹R¹² or nitro; C₂-C₁₀ alkyl substituted with cyano; C₁-C₁₀ alkoxy; C₁-C₁₀ haloalkoxy; C₁-C₁₀ alkylthio; C₁-C₁₀ haloalkylthio; NR¹²R¹⁷; or phenyl, pyridyl, furanyl, thienyl, naphthyl, benzofuranyl, benzothienyl, or quinolinyl each optionally substituted with R¹⁴, R¹⁵, and R¹⁶;

R⁸ is hydrogen; C₁-C₄ alkyl; or -C(=O)R¹⁰;


R⁹ is hydrogen; C₂-C₁₀ alkyl; C₃-C₇ cycloalkyl; C₃-C₁₀ alkenyl; C₃-C₁₀ alkynyl; C₃-C₁₀ haloalkyl; C₃-C₁₀ haloalkenyl; C₃-C₁₀ haloalkynyl; C₃-C₁₀ alkoxyalkyl other than butoxyethyl; C₂-C₁₀ alkylthioalkyl; C₂-C₁₀ alkylsulfinylalkyl; C₂-C₁₀ alkylsulfonylalkyl; C₄-C₁₀ cycloalkylalkyl; C₄-C₁₀ alkenyloxyalkyl; C₄-C₁₀ alkynyloxyalkyl; C₄-C₁₀ (cycloalkyl)oxyalkyl; C₄-C₁₀ alkenylthioalkyl; C₄-C₁₀ alkynylthioalkyl; C₆-C₁₀ (cycloalkyl)thioalkyl; C₂-C₁₀ haloalkoxyalkyl; C₄-C₁₀ haloalkenyloxyalkyl; C₄-C₁₀ haloalkynyloxyalkyl; C₄-C₁₀ alkoxyalkenyl; C₄-C₁₀ alkoxyalkynyl; C₄-C₁₀ alkylthioalkenyl; C₄-C₁₀ alkylthioalkynyl; C₄-C₁₀ trialkylsilylalkyl;

- C_1 - C_{10} alkyl substituted with $NR^{11}R^{12}$; C_4 - C_{10} cyanoalkyl; C_2 - C_{10} nitroalkyl; C_1 - C_8 alkyl substituted with CO_2R^{11} ; pyridyl, furanyl, thienyl, or naphthyl each optionally substituted with R^{14} , R^{15} , and R^{16} ; $-N=CR^{11}R^{11}$; $-NR^{12}R^{17}$; $-OR^{12}$; or $-NC(=Q)NR^{11}R^{12}$; or R^3 and R^4 are both iodine and R^9 is phenyl optionally substituted with R^{14} , R^{15} , and R^{16} ; or R^7 and R^9 are taken together to form $-CH_2(CH_2)_mCH_2-$; R^{10} is hydrogen; C_1 - C_4 alkyl; C_1 - C_4 alkoxy; or $NR^{11}R^{12}$; R^{11} is independently hydrogen; C_1 - C_4 alkyl; or phenyl optionally substituted with at least one R^{13} ; R^{12} is independently hydrogen; C_1 - C_8 alkyl; or phenyl optionally substituted with at least one R^{13} ; or R^{11} and R^{12} are taken together to form $-CH_2CH_2CH_2CH_2-$, $-CH_2(CH_2)_3CH_2-$, $-CH_2CH_2OCH_2CH_2-$, $-CH_2CH(Me)CH_2CH(Me)CH_2-$, or $-CH_2CH(Me)OCH(Me)CH_2-$; R^{13} is independently halogen; C_1 - C_4 alkyl; C_1 - C_4 alkoxy; C_1 - C_4 haloalkyl; nitro; or cyano; R^{14} is independently C_1 - C_6 alkyl; C_1 - C_6 alkoxy; C_1 - C_6 haloalkyl; halogen; C_2 - C_8 alkynyl; C_1 - C_6 thioalkyl; phenyl or phenoxy each optionally substituted with at least one R^{13} ; cyano; nitro; C_1 - C_6 haloalkoxy; C_1 - C_6 haloalkylthio; C_2 - C_6 alkenyl; C_2 - C_6 haloalkenyl; acetyl; CO_2Me ; or $N(C_1-C_2\text{ alkyl})_2$; R^{15} is independently methyl; ethyl; methoxy; methylthio; halogen; or trifluoromethyl; R^{16} is independently halogen; and R^{17} is independently C_1 - C_8 alkyl; or phenyl optionally substituted with at least one R^{13} ; or an *N*-oxide or agriculturally-suitable salt thereof; and at least one of (a) a surfactant, (b) an organic solvent, and (c) at least one solid or liquid diluent.
15. A method for controlling wheat powdery mildew comprising applying to the plant or portion thereof to be protected, to the media in which the plant to be protected is growing, or to the plant seed or seedling to be protected an effective amount of a compound of Claim 1.

AMENDED CLAIMS

[received by the International Bureau on 31 October 1994 (31.10.94); original claim 1 amended; new claims 2 and 14-17 added; original claims 2-12 and 13-15; renumbered as claims 3-13 and 18-20 respectively (15 pages)]

1. A compound of Formula I, II, or III



wherein:

n is 0, 1 or 2;

Q is independently O or S;

10 R¹ is C₃-C₁₀ alkyl; C₃-C₅ cycloalkyl; C₄-C₁₀ alkenyl; C₄-C₁₀ alkynyl; C₁-C₁₀ haloalkyl; C₃-C₁₀ haloalkenyl; C₃-C₁₀ haloalkynyl; C₂-C₁₀ alkoxyalkyl; C₂-C₁₀ alkylthioalkyl; C₂-C₁₀ alkylsulfinylalkyl; C₂-C₁₀ alkylsulfonylalkyl; cyclopropylmethyl; C₅-C₁₀ cycloalkylalkyl; C₄-C₁₀ alkenyloxyalkyl; C₄-C₁₀ alkynyloxyalkyl; C₄-C₁₀ (cycloalkyl)oxyalkyl; C₄-C₁₀ alkenylthioalkyl; C₄-C₁₀ alkynylthioalkyl; C₆-C₁₀ (cycloalkyl)thioalkyl; C₂-C₁₀ haloalkoxyalkyl; C₄-C₁₀ haloalkenyloxyalkyl; C₄-C₁₀ haloalkynyloxyalkyl; C₄-C₁₀ alkoxyalkenyl; C₄-C₁₀ alkoxyalkynyl; C₄-C₁₀ alkylthioalkenyl; C₄-C₁₀ alkylthioalkynyl; C₄-C₁₀ trialkylsilylalkyl; C₁-C₁₀ alkyl substituted with NR¹¹R¹², nitro, cyano, or phenyl optionally substituted with R¹⁴, R¹⁵, and R¹⁶; C₁-C₁₀ alkoxy; C₁-C₁₀ haloalkoxy; C₁-C₁₀ alkylthio; C₁-C₁₀ haloalkylthio; NR¹¹R¹²; or pyridyl, furanyl, thienyl, naphthyl, benzofuranyl, benzothienyl, or quinoliny each optionally substituted with R¹⁴, R¹⁵, and R¹⁶;

25 R² is C₃-C₁₀ alkyl; C₆-C₇ cycloalkyl; C₃-C₁₀ alkenyl; C₃-C₁₀ alkynyl; C₁-C₁₀ haloalkyl; C₃-C₁₀ haloalkenyl; C₃-C₁₀ haloalkynyl; C₂-C₁₀ alkoxyalkyl; C₂-C₁₀ alkylthioalkyl; C₂-C₁₀ alkylsulfinylalkyl; C₂-C₁₀ alkylsulfonylalkyl; C₄-C₁₀ cycloalkylalkyl; C₄-C₁₀ alkenyloxyalkyl; C₄-C₁₀ alkynyloxyalkyl; C₄-C₁₀ (cycloalkyl)oxyalkyl; C₄-C₁₀ alkenylthioalkyl; C₄-C₁₀ alkynylthioalkyl; C₆-C₁₀ (cycloalkyl)thioalkyl; C₂-C₁₀ haloalkoxyalkyl; C₄-C₁₀ haloalkenyloxyalkyl; C₄-C₁₀ haloalkynyloxyalkyl; C₄-C₁₀ alkoxyalkenyl; C₄-C₁₀ alkoxyalkynyl; C₄-C₁₀ alkylthioalkenyl; C₄-C₁₀ alkylthioalkynyl; C₄-C₁₀ trialkylsilylalkyl; C₃-C₁₀ cyanoalkyl; C₂-C₁₀ nitroalkyl; C₁-C₁₀ alkyl substituted with CO₂R¹¹, NR¹¹R¹², or phenyl

30

optionally substituted with R^{13} , R^{15} , and R^{16} ; phenyl optionally substituted with R^{13} , R^{15} , and R^{16} ; $-N=CR^{11}R^{11}$; or $-NR^{11}R^{12}$; or

R^1 and R^2 are taken together to form $-\text{CH}_2(\text{CH}_2)_m\text{CH}_2-$;

m is 1-4;

- 5 R^3 is halogen; C_1 - C_8 alkyl; C_3 - C_8 cycloalkyl; C_2 - C_8 alkenyl; C_2 - C_8 alkynyl; C_1 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; C_3 - C_8 haloalkynyl; C_1 - C_8 alkoxy; C_1 - C_8 haloalkoxy; C_3 - C_8 alkenyloxy; C_3 - C_8 alkynyloxy; C_1 - C_8 alkylthio; C_3 - C_8 alkenylthio; C_3 - C_8 alkynylthio; C_1 - C_8 alkylsulfinyl; C_1 - C_8 alkylsulfonyl; C_2 - C_8 alkoxyalkyl; C_2 - C_8 alkylthioalkyl; C_2 - C_8 alkylsulfinylalkyl; C_2 - C_8 alkylsulfonylalkyl; C_4 - C_8 cycloalkylalkyl; C_3 - C_8 trialkylsilyl; nitro; $NR^{11}R^{12}$; C_5 - C_8 trialkylsilylalkynyl; or phenyl optionally substituted with at least one R^{13} ;
- 10 R^4 is hydrogen; halogen; C_1 - C_4 alkyl; C_1 - C_4 haloalkyl; C_1 - C_4 alkoxy; or C_1 - C_4 haloalkoxy;
- 15 R^5 is C_3 - C_5 alkyl; C_7 - C_{10} alkyl; C_4 - C_7 alkenyl; C_3 - C_5 alkynyl; C_1 - C_{10} haloalkyl; C_5 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_2 - C_{10} alkoxyalkyl other than methoxypropyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_4 - C_{10} cycloalkylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_1 - C_{10} alkyl substituted with $NR^{11}R^{12}$, nitro, or phenyl optionally substituted with at least one of R^{14} , R^{15} , and R^{16} ; C_2 - C_{10} alkyl substituted with cyano; C_1 - C_{10} alkoxy; C_1 - C_{10} haloalkoxy; C_1 - C_{10} alkylthio; C_1 - C_{10} haloalkylthio; $NR^{11}R^{12}$; or phenyl, furanyl, thienyl, naphthyl, benzofuranyl, or benzothienyl each optionally substituted with R^{14} , R^{15} , and R^{16} ;
- 20 R^6 is C_3 - C_{10} alkyl; C_3 - C_7 alkenyl; C_3 - C_{10} alkynyl; C_1 - C_{10} haloalkyl; C_3 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_2 - C_{10} alkoxyalkyl other than propoxymethyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_4 - C_{10} cycloalkylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_5 - C_{10} cyanoalkyl; C_2 - C_{10} nitroalkyl; or C_3 - C_{10} alkyl substituted with CO_2R^{11} , $NR^{11}R^{12}$, or phenyl
- 30
- 35

optionally substituted with R^{13} , R^{15} , and R^{16} ; or phenyl optionally substituted with R^{13} , R^{15} , and R^{16} ; or

R^5 and R^6 are taken together to form $-\text{CH}_2(\text{CH}_2)_m\text{CH}_2-$;

R^7 is C_3 - C_{10} alkyl; C_3 - C_7 cycloalkyl; C_4 - C_7 alkenyl; propynyl; C_5 - C_{10} alkynyl; C_2 - C_{10} haloalkyl; C_3 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_2 - C_{10} alkoxyalkyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_1 - C_{10} alkyl substituted with $\text{NR}^{11}\text{R}^{12}$ or nitro; C_2 - C_{10} alkyl substituted with cyano; C_1 - C_{10} alkoxy; C_1 - C_{10} haloalkoxy; C_1 - C_{10} alkylthio; C_1 - C_{10} haloalkylthio; $\text{NR}^{12}\text{R}^{17}$; or phenyl, pyridyl, furanyl, thienyl, naphthyl, benzofuranyl, benzothienyl, or quinolinyl each optionally substituted with R^{14} , R^{15} , and R^{16} ;

R^8 is hydrogen; C_1 - C_4 alkyl; or $-\text{C}(=\text{O})\text{R}^{10}$;

R^9 is hydrogen; C_2 - C_{10} alkyl; C_3 - C_7 cycloalkyl; C_3 - C_{10} alkenyl; C_3 - C_{10} alkynyl; C_3 - C_{10} haloalkyl; C_3 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_3 - C_{10} alkoxyalkyl other than butoxyethyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_4 - C_{10} cycloalkylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_1 - C_{10} alkyl substituted with $\text{NR}^{11}\text{R}^{12}$; C_4 - C_{10} cyanoalkyl; C_2 - C_{10} nitroalkyl; C_1 - C_8 alkyl substituted with CO_2R^{11} ; pyridyl, furanyl, thienyl, or naphthyl each optionally substituted with R^{14} , R^{15} , and R^{16} ; $-\text{N}=\text{CR}^{11}\text{R}^{11}$; $-\text{NR}^{12}\text{R}^{17}$; $-\text{OR}^{12}$; or $-\text{NC}(=\text{Q})\text{NR}^{11}\text{R}^{12}$; or R^3 and R^4 are both iodine and R^9 is phenyl optionally substituted with R^{14} , R^{15} , and R^{16} ; or

R^7 and R^9 are taken together to form $-\text{CH}_2(\text{CH}_2)_m\text{CH}_2-$;

R^{10} is hydrogen; C_1 - C_4 alkyl; C_1 - C_4 alkoxy; or $\text{NR}^{11}\text{R}^{12}$;

R^{11} is independently hydrogen; C_1 - C_4 alkyl; or phenyl optionally substituted with at least one R^{13} ;

R^{12} is independently hydrogen; C_1 - C_8 alkyl; or phenyl optionally substituted with at least one R^{13} ; or

R^{11} and R^{12} are taken together to form $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2(\text{CH}_2)_3\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}(\text{Me})\text{CH}_2\text{CH}(\text{Me})\text{CH}_2-$, or $-\text{CH}_2\text{CH}(\text{Me})\text{OCH}(\text{Me})\text{CH}_2-$;

R^{13} is independently halogen; C_1 - C_4 alkyl; C_1 - C_4 alkoxy; C_1 - C_4 haloalkyl; nitro; or cyano;

R^{14} is independently C_1 - C_6 alkyl; C_1 - C_6 alkoxy; C_1 - C_6 haloalkyl; halogen; C_2 - C_8 alkynyl; C_1 - C_6 thioalkyl; phenyl or phenoxy each optionally substituted with at least one R^{13} ; cyano; nitro; C_1 - C_6 haloalkoxy; C_1 - C_6 haloalkylthio; C_2 - C_6 alkenyl; C_2 - C_6 haloalkenyl; acetyl; CO_2Me ; or $\text{N}(\text{C}_1\text{-C}_2 \text{ alkyl})_2$;

R^{15} is independently methyl; ethyl; methoxy; methylthio; halogen; or trifluoromethyl;

R^{16} is independently halogen; and

R^{17} is independently C_1 - C_8 alkyl; or phenyl optionally substituted with at least one R^{13} ; or

an *N*-oxide or agriculturally-suitable salt thereof; provided that when R^1 is cyclopropylmethyl, R^2 is $\text{CH}_2\text{CH}_2\text{CH}_3$, R^3 is 6-Br and R^4 is H.

2. A compound of Formula I of Claim 1 wherein:

R^1 is C_3 - C_{10} alkyl; C_3 - C_5 cycloalkyl; C_4 - C_{10} alkenyl; C_4 - C_{10} alkynyl; C_1 - C_{10} haloalkyl; C_3 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_2 - C_{10} alkoxyalkyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_5 - C_{10} cycloalkylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_1 - C_{10} alkyl substituted with $\text{NR}^{11}\text{R}^{12}$, nitro, cyano, or phenyl optionally substituted with R^{14} , R^{15} , and R^{16} ; C_1 - C_{10} alkoxy; C_1 - C_{10} haloalkoxy; C_1 - C_{10} alkylthio; C_1 - C_{10} haloalkylthio; $\text{NR}^{11}\text{R}^{12}$; or pyridyl, furanyl, thienyl, naphthyl, benzofuranyl, benzothienyl, or quinolinyl each optionally substituted with R^{14} , R^{15} , and R^{16} .

3. A compound of Claim 2 wherein:

Q is O;

R^1 is C_3 - C_8 alkyl; C_4 - C_8 alkenyl; C_4 - C_8 alkynyl; C_1 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; C_2 - C_8 alkoxyalkyl; C_2 - C_8 alkylthioalkyl; C_5 - C_8 cycloalkylalkyl; C_2 - C_8 alkyl substituted with cyano; C_1 - C_8 alkoxy; C_1 - C_8 haloalkoxy; C_1 - C_8

alkylthio; or C₄-C₈ alkenyloxyalkyl; or pyridyl, furanyl, or thienyl each optionally substituted with R¹⁴ and R¹⁵;

R² is C₃-C₈ alkyl; C₃-C₈ alkenyl; C₃-C₈ alkynyl; C₁-C₈ haloalkyl; C₃-C₈ haloalkenyl; C₃-C₈ alkoxyalkyl; C₂-C₈ alkylthioalkyl; C₄-C₈ cycloalkylalkyl; C₃-C₈ cyanoalkyl; C₄-C₈ alkenyloxyalkyl; or phenyl optionally substituted with R¹³;

R³ is halogen; C₁-C₈ alkyl; C₂-C₈ alkynyl; C₃-C₈ cycloalkyl; C₁-C₈ haloalkyl; C₁-C₈ alkoxy; C₁-C₈ haloalkoxy; C₁-C₈ alkylthio; C₁-C₈ alkylsulfonyl; C₂-C₈ alkoxyalkyl; C₂-C₈ alkylthioalkyl; C₄-C₈ cycloalkylalkyl; or C₅-C₈ trialkylsilylalkynyl; and

R¹⁴ is methyl; ethyl; methoxy; ethoxy; C₁-C₂ haloalkyl; halogen; acetylenyl; propargyl; methylthio; ethylthio; cyano; nitro; C₁-C₂ haloalkoxy; vinyl; allyl; acetyl; CO₂Me; or N(C₁-C₂ alkyl)₂.

4. A compound of Formula II of Claim 1 wherein:

Q is O;

n is 0;

R³ is halogen; C₁-C₈ alkyl; C₂-C₈ alkynyl; C₃-C₈ cycloalkyl; C₁-C₈ haloalkyl; C₁-C₈ alkoxy; C₁-C₈ haloalkoxy; C₁-C₈ alkylthio; C₁-C₈ alkylsulfonyl; C₂-C₈ alkoxyalkyl; C₂-C₈ alkylthioalkyl; C₄-C₈ cycloalkylalkyl; or C₅-C₈ trialkylsilylalkynyl;

R⁵ is C₃-C₅ alkyl; C₄-C₇ alkenyl; C₃-C₅ alkynyl; C₁-C₈ haloalkyl; C₅-C₈ haloalkenyl; C₂-C₈ alkoxyalkyl other than methoxypropyl; C₂-C₈ alkylthioalkyl; C₄-C₈ cycloalkylalkyl; C₂-C₈ alkyl substituted with cyano; C₁-C₈ alkoxy; C₁-C₈ haloalkoxy; C₁-C₈ alkylthio; or C₄-C₈ alkenyloxyalkyl; or phenyl, furanyl, or thienyl each optionally substituted with R¹⁴ and R¹⁵;

R⁶ is C₃-C₈ alkyl; C₃-C₇ alkenyl; C₃-C₈ alkynyl; C₁-C₈ haloalkyl; C₃-C₈ haloalkenyl; C₃-C₈ alkoxyalkyl other than propoxymethyl; C₂-C₈ alkylthioalkyl; C₄-C₈ cycloalkylalkyl; C₅-C₈ cyanoalkyl; C₄-C₈ alkenyloxyalkyl; phenyl optionally substituted with R¹³; or C₃-C₅ alkyl substituted with phenyl optionally substituted with R¹³ and R¹⁵; and

R¹⁴ is methyl; ethyl; methoxy; ethoxy; C₁-C₂ haloalkyl; halogen; acetylenyl; propargyl; methylthio; ethylthio; cyano; nitro; C₁-C₂ haloalkoxy; vinyl; allyl; acetyl; CO₂Me; or N(C₁-C₂ alkyl)₂.

5. A compound of Formula III of Claim 1 wherein:

Q is O;

- R^3 is halogen; C_1 - C_8 alkyl; C_2 - C_8 alkynyl; C_3 - C_8 cycloalkyl; C_1 - C_8 haloalkyl; C_1 - C_8 alkoxy; C_1 - C_8 haloalkoxy; C_1 - C_8 alkylthio; C_1 - C_8 alkylsulfonyl; C_2 - C_8 alkoxyalkyl; C_2 - C_8 alkylthioalkyl; C_4 - C_8 cycloalkylalkyl; or C_5 - C_8 trialkylsilylalkynyl;
- 5 R^7 is C_3 - C_8 alkyl; C_4 - C_7 alkenyl; propynyl; C_2 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; C_2 - C_8 alkoxyalkyl; C_2 - C_8 alkylthioalkyl; C_2 - C_8 alkyl substituted with cyano; C_1 - C_8 alkoxy; C_1 - C_8 haloalkoxy; C_1 - C_8 alkylthio; or C_4 - C_8 alkenyloxyalkyl; or phenyl, pyridyl, furanyl, or thienyl each optionally substituted with R^{14} and R^{15} ;
- 10 R^9 is C_3 - C_8 alkyl; C_3 - C_8 alkenyl; C_3 - C_8 alkynyl; C_3 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; C_3 - C_8 alkoxyalkyl; C_2 - C_8 alkylthioalkyl; C_4 - C_8 cycloalkylalkyl; C_4 - C_8 cyanoalkyl; C_4 - C_8 alkenyloxyalkyl; $-NR^{12}R^{17}$; or R^3 and R^4 are both iodine and R^9 is phenyl optionally substituted with R^{14} and R^{15} ; and
- 15 R^{14} is methyl; ethyl; methoxy; ethoxy; C_1 - C_2 haloalkyl; halogen; acetylenyl; propargyl; methylthio; ethylthio; cyano; nitro; C_1 - C_2 haloalkoxy; vinyl; allyl; acetyl; CO_2Me ; or $N(C_1-C_2 \text{ alkyl})_2$.
6. A compound of Claim 3 wherein:
- 20 R^1 is C_3 - C_8 alkyl; C_4 - C_8 alkenyl; C_4 - C_8 alkynyl; C_3 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; C_3 - C_8 alkoxyalkyl; or thienyl optionally substituted with at least one of R^{14} and R^{15} ;
- R^2 is C_3 - C_8 alkyl; C_3 - C_8 alkenyl; C_3 - C_8 alkynyl; C_3 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; C_3 - C_8 alkoxyalkyl; or phenyl optionally substituted with R^{13} ;
- 25 R^3 is halogen; C_1 - C_4 alkyl; C_1 - C_4 haloalkyl; C_1 - C_4 alkoxy; C_1 - C_4 haloalkoxy; acetylenyl; or trimethylsilylacetylenyl; and
- R^{14} is methyl; ethyl; methoxy; methylthio; halogen; trifluoromethyl; or $N(C_1-C_2 \text{ alkyl})_2$.
7. A compound of Claim 4 wherein:
- 30 R^3 is halogen; C_1 - C_4 alkyl; C_1 - C_4 haloalkyl; C_1 - C_4 alkoxy; C_1 - C_4 haloalkoxy; acetylenyl; or trimethylsilylacetylenyl;
- R^5 is C_3 - C_5 alkyl; C_4 - C_7 alkenyl; C_3 - C_5 alkynyl; C_3 - C_8 haloalkyl; C_5 - C_8 haloalkenyl; C_3 - C_8 alkoxyalkyl; or phenyl or thienyl each optionally substituted with R^{14} and R^{15} ;
- 35 R^6 is C_3 - C_8 alkyl; C_3 - C_7 alkenyl; C_3 - C_8 alkynyl; C_3 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; C_3 - C_8 alkoxyalkyl; or phenyl optionally substituted with R^{13} ; and

R¹⁴ is methyl; ethyl; methoxy; methylthio; halogen; trifluoromethyl; or
N(C₁-C₂ alkyl)₂.

8. A compound of Claim 5 wherein:

5 R³ is halogen; C₁-C₄ alkyl; C₁-C₄ haloalkyl; C₁-C₄ alkoxy; C₁-C₄ haloalkoxy;
acetylenyl; or trimethylsilylacetylenyl;

R⁷ is C₃-C₈ alkyl; C₄-C₇ alkenyl; propynyl; C₃-C₈ haloalkyl; C₃-C₈ haloalkenyl;
C₃-C₈ alkoxy; C₃-C₈ alkoxyalkyl; or phenyl or thienyl each optionally
substituted with R¹⁴ and R¹⁵;

10 R⁹ is C₃-C₈ alkyl; C₃-C₈ alkenyl; C₃-C₈ alkynyl; C₃-C₈ haloalkyl; C₃-C₈
haloalkenyl; C₃-C₈ alkoxyalkyl; -NR¹²R¹⁷; or R³ and R⁴ are both iodine and
R⁹ is phenyl optionally substituted with R¹⁴ and R¹⁵; and

R¹⁴ is methyl; ethyl; methoxy; methylthio; halogen; trifluoromethyl; or
N(C₁-C₂ alkyl)₂.

15

9. A compound of Claim 6 wherein:

R¹ is C₃-C₈ alkyl; C₄-C₈ alkenyl; C₄-C₈ alkynyl; C₃-C₈ haloalkyl; or C₃-C₈
haloalkenyl;

20 R² is C₃-C₈ alkyl; C₃-C₈ alkenyl; C₃-C₈ alkynyl; C₃-C₈ haloalkyl; C₃-C₈
haloalkenyl; or phenyl optionally substituted with R¹³;

R³ is halogen; and

R⁴ is hydrogen or halogen.

10. A compound of Claim 7 wherein:

25 R³ is halogen;

R⁴ is hydrogen or halogen;

R⁵ is C₃-C₅ alkyl; C₄-C₇ alkenyl; C₃-C₅ alkynyl; C₃-C₈ haloalkyl; or C₅-C₈
haloalkenyl; or phenyl optionally substituted with R¹⁴ and R¹⁵; and

30 R⁶ is C₃-C₈ alkyl; C₃-C₇ alkenyl; C₃-C₈ alkynyl; C₃-C₈ haloalkyl; C₃-C₈
haloalkenyl; or phenyl optionally substituted with R¹³.

11. A compound of Claim 8 wherein:

R³ is halogen;




R⁴ is hydrogen or halogen;

35 R⁷ is C₃-C₈ alkyl; C₄-C₇ alkenyl; propynyl; C₃-C₈ haloalkyl; or C₃-C₈ haloalkenyl;
or phenyl optionally substituted with R¹⁴ and R¹⁵; and

R⁹ is C₃-C₈ alkyl; C₃-C₈ alkenyl; C₃-C₈ alkynyl; C₃-C₈ haloalkyl; C₃-C₈ haloalkenyl; -NR¹²R¹⁷; or R³ and R⁴ are both iodine and R⁹ is phenyl optionally substituted with R¹⁴ and R¹⁵.

- 5 12. A compound of Claim 9 which is selected from:
6-bromo-3-propyl-2-propyloxy-4(3H)-quinazolinone;
6,8-diiodo-3-propyl-2-propyloxy-4(3H)-quinazolinone; and
6-iodo-3-propyl-2-propyloxy-4(3H)-quinazolinone.
- 10 13. A compound of Claim 11 which is
6,8-diiodo-3-propyl-2-(phenylamino)-4(3H)-quinazolinone.
- 15 14. A compound of Claim 1 which is
6-bromo-3-(cyclopropylmethyl)-2-propyloxy-4(3H)-quinazolinone.
- 20 15. A method for controlling plant diseases caused by fungal plant pathogens,
comprising applying to the plant or portion thereof to be protected, to the media in
which the plant to be protected is growing, or to the plant seed or seedling to be
protected an effective amount of the compound of Claim 14.
- 25 16. A fungicidal composition comprising an effective amount of the compound
of Claim 14 and at least one of (a) a surfactant, (b) an organic solvent, and (c) at least
one solid or liquid diluent.
- 30 17. A method for controlling wheat powdery mildew comprising applying to the
plant or portion thereof to be protected, to the media in which the plant to be protected
is growing, or to the plant seed or seedling to be protected an effective amount of the
compound of Claim 14.
- 30 18. A method for controlling plant diseases caused by fungal plant pathogens
comprising applying to the plant or portion thereof to be protected, to the media in
which the plant to be protected is growing, or to the plant seed or seedling to be
protected an effective amount of a compound of Formula I, II, or III:

64



wherein:

n is 0, 1 or 2;

5 Q is independently O or S;

R¹ is C₃-C₁₀ alkyl; C₃-C₅ cycloalkyl; C₄-C₁₀ alkenyl; C₄-C₁₀ alkynyl; C₁-C₁₀ haloalkyl; C₃-C₁₀ haloalkenyl; C₃-C₁₀ haloalkynyl; C₂-C₁₀ alkoxyalkyl; C₂-C₁₀ alkylthioalkyl; C₂-C₁₀ alkylsulfinylalkyl; C₂-C₁₀ alkylsulfonylalkyl; C₅-C₁₀ cycloalkylalkyl; C₄-C₁₀ alkenyloxyalkyl; C₄-C₁₀ alkynyloxyalkyl; C₄-C₁₀ (cycloalkyl)oxyalkyl; C₄-C₁₀ alkenylthioalkyl; C₄-C₁₀ alkynylthioalkyl; C₆-C₁₀ (cycloalkyl)thioalkyl; C₂-C₁₀ haloalkoxyalkyl; C₄-C₁₀ haloalkenyloxyalkyl; C₄-C₁₀ haloalkynyloxyalkyl; C₄-C₁₀ alkoxyalkenyl; C₄-C₁₀ alkoxyalkynyl; C₄-C₁₀ alkylthioalkenyl; C₄-C₁₀ alkylthioalkynyl; C₄-C₁₀ trialkylsilylalkyl; C₁-C₁₀ alkyl substituted with NR¹¹R¹², nitro, cyano, or phenyl optionally substituted with R¹⁴, R¹⁵, and R¹⁶; C₁-C₁₀ alkoxy; C₁-C₁₀ haloalkoxy; C₁-C₁₀ alkylthio; C₁-C₁₀ haloalkylthio; NR¹¹R¹²; or pyridyl, furanyl, thienyl, naphthyl, benzofuranyl, benzothienyl, or quinolinyl each optionally substituted with R¹⁴, R¹⁵, and R¹⁶;

20 R² is C₃-C₁₀ alkyl; C₆-C₇ cycloalkyl; C₃-C₁₀ alkenyl; C₃-C₁₀ alkynyl; C₁-C₁₀ haloalkyl; C₃-C₁₀ haloalkenyl; C₃-C₁₀ haloalkynyl; C₂-C₁₀ alkoxyalkyl; C₂-C₁₀ alkylthioalkyl; C₂-C₁₀ alkylsulfinylalkyl; C₂-C₁₀ alkylsulfonylalkyl; C₄-C₁₀ cycloalkylalkyl; C₄-C₁₀ alkenyloxyalkyl; C₄-C₁₀ alkynyloxyalkyl; C₄-C₁₀ (cycloalkyl)oxyalkyl; C₄-C₁₀ alkenylthioalkyl; C₄-C₁₀ alkynylthioalkyl; C₆-C₁₀ (cycloalkyl)thioalkyl; C₂-C₁₀ haloalkoxyalkyl; C₄-C₁₀ haloalkenyloxyalkyl; C₄-C₁₀ haloalkynyloxyalkyl; C₄-C₁₀ alkoxyalkenyl; C₄-C₁₀ alkoxyalkynyl; C₄-C₁₀ alkylthioalkenyl; C₄-C₁₀ alkylthioalkynyl; C₄-C₁₀ trialkylsilylalkyl; C₃-C₁₀ cyanoalkyl; C₂-C₁₀ nitroalkyl; C₁-C₁₀ alkyl substituted with CO₂R¹¹, NR¹¹R¹², or phenyl optionally substituted with R¹³, R¹⁵, and R¹⁶; phenyl optionally substituted with R¹³, R¹⁵, and R¹⁶; -N=CR¹¹R¹¹; or -NR¹¹R¹²; or

R¹ and R² are taken together to form -CH₂(CH₂)_mCH₂-;

m is 1-4;

- 5 R^3 is halogen; C_1 - C_8 alkyl; C_3 - C_8 cycloalkyl; C_2 - C_8 alkenyl; C_2 - C_8 alkynyl; C_1 - C_8 haloalkyl; C_3 - C_8 haloalkenyl; C_3 - C_8 haloalkynyl; C_1 - C_8 alkoxy; C_1 - C_8 haloalkoxy; C_3 - C_8 alkenyloxy; C_3 - C_8 alkynyloxy; C_1 - C_8 alkylthio; C_3 - C_8 alkenylthio; C_3 - C_8 alkynylthio; C_1 - C_8 alkylsulfinyl; C_1 - C_8 alkylsulfonyl; C_2 - C_8 alkoxyalkyl; C_2 - C_8 alkylthioalkyl; C_2 - C_8 alkylsulfinylalkyl; C_2 - C_8 alkylsulfonylalkyl; C_4 - C_8 cycloalkylalkyl; C_3 - C_8 trialkylsilyl; nitro; $NR^{11}R^{12}$; C_5 - C_8 trialkylsilylalkynyl; or phenyl optionally substituted with at least one R^{13} ;
- 10 R^4 is hydrogen; halogen; C_1 - C_4 alkyl; C_1 - C_4 haloalkyl; C_1 - C_4 alkoxy; or C_1 - C_4 haloalkoxy;
- 15 R^5 is C_3 - C_5 alkyl; C_7 - C_{10} alkyl; C_4 - C_7 alkenyl; C_3 - C_5 alkynyl; C_1 - C_{10} haloalkyl; C_5 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_2 - C_{10} alkoxyalkyl other than methoxypropyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_4 - C_{10} cycloalkylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_1 - C_{10} alkyl substituted with $NR^{11}R^{12}$, nitro, or phenyl optionally substituted with at least one of R^{14} , R^{15} , and R^{16} ; C_2 - C_{10} alkyl substituted with cyano; C_1 - C_{10} alkoxy; C_1 - C_{10} haloalkoxy; C_1 - C_{10} alkylthio; C_1 - C_{10} haloalkylthio; $NR^{11}R^{12}$; or phenyl, furanyl, thienyl, naphthyl, benzofuranyl, or benzothienyl each optionally substituted with R^{14} , R^{15} , and R^{16} ;
- 20 R^6 is C_3 - C_{10} alkyl; C_3 - C_7 alkenyl; C_3 - C_{10} alkynyl; C_1 - C_{10} haloalkyl; C_3 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_3 - C_{10} alkoxyalkyl other than propoxymethyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_4 - C_{10} cycloalkylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_5 - C_{10} cyanoalkyl; C_2 - C_{10} nitroalkyl; or C_3 - C_{10} alkyl substituted with CO_2R^{11} , $NR^{11}R^{12}$, or phenyl optionally substituted with R^{13} , R^{15} , and R^{16} ; or phenyl optionally substituted with R^{13} , R^{15} , and R^{16} ; or
- 30 R^5 and R^6 are taken together to form $-CH_2(CH_2)_mCH_2-$;
- 35

- 5 R^7 is C_3 - C_{10} alkyl; C_3 - C_7 cycloalkyl; C_4 - C_7 alkenyl; propynyl; C_5 - C_{10} alkynyl; C_2 - C_{10} haloalkyl; C_3 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_2 - C_{10} alkoxyalkyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_1 - C_{10} alkyl substituted with $NR^{11}R^{12}$ or nitro; C_2 - C_{10} alkyl substituted with cyano; C_1 - C_{10} alkoxy; C_1 - C_{10} haloalkoxy; C_1 - C_{10} alkylthio; C_1 - C_{10} haloalkylthio; $NR^{12}R^{17}$; or phenyl, pyridyl, furanyl, thienyl, naphthyl, benzofuranyl, benzothienyl, or quinolynyl each optionally substituted with R^{14} , R^{15} , and R^{16} ;
- 10 R^8 is hydrogen; C_1 - C_4 alkyl; or $-C(=O)R^{10}$;
- 15 R^9 is hydrogen; C_2 - C_{10} alkyl; C_3 - C_7 cycloalkyl; C_3 - C_{10} alkenyl; C_3 - C_{10} alkynyl; C_3 - C_{10} haloalkyl; C_3 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_3 - C_{10} alkoxyalkyl other than butoxyethyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_4 - C_{10} cycloalkylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_1 - C_{10} alkyl substituted with $NR^{11}R^{12}$; C_4 - C_{10} cyanoalkyl; C_2 - C_{10} nitroalkyl; C_1 - C_8 alkyl substituted with CO_2R^{11} ; pyridyl, furanyl, thienyl, or naphthyl each optionally substituted with R^{14} , R^{15} , and R^{16} ; $-N=CR^{11}R^{11}$; $-NR^{12}R^{17}$; $-OR^{12}$; or $-NC(=Q)NR^{11}R^{12}$; or R^3 and R^4 are both iodine and R^9 is phenyl optionally substituted with R^{14} , R^{15} , and R^{16} ; or
- 20 R^7 and R^9 are taken together to form $-CH_2(CH_2)_mCH_2-$;
- 25 R^{10} is hydrogen; C_1 - C_4 alkyl; C_1 - C_4 alkoxy; or $NR^{11}R^{12}$;
- 30 R^{11} is independently hydrogen; C_1 - C_4 alkyl; or phenyl optionally substituted with at least one R^{13} ;
- 35 R^{12} is independently hydrogen; C_1 - C_8 alkyl; or phenyl optionally substituted with at least one R^{13} ; or
- R^{11} and R^{12} are taken together to form $-CH_2CH_2CH_2CH_2-$, $-CH_2(CH_2)_3CH_2-$, $-CH_2CH_2OCH_2CH_2-$, $-CH_2CH(Me)CH_2CH(Me)CH_2-$, or $-CH_2CH(Me)OCH(Me)CH_2-$;

R^{13} is independently halogen; C_1 - C_4 alkyl; C_1 - C_4 alkoxy; C_1 - C_4 haloalkyl; nitro; or cyano;

R^{14} is independently C_1 - C_6 alkyl; C_1 - C_6 alkoxy; C_1 - C_6 haloalkyl; halogen; C_2 - C_8 alkynyl; C_1 - C_6 thioalkyl; phenyl or phenoxy each optionally substituted with at least one R^{13} ; cyano; nitro; C_1 - C_6 haloalkoxy; C_1 - C_6 haloalkylthio; C_2 - C_6 alkenyl; C_2 - C_6 haloalkenyl; acetyl; CO_2Me ; or $N(C_1$ - C_2 alkyl) $_2$;




R^{15} is independently methyl; ethyl; methoxy; methylthio; halogen; or trifluoromethyl;

R^{16} is independently halogen; and

R^{17} is independently C_1 - C_8 alkyl; or phenyl optionally substituted with at least one R^{13} ; or

an *N*-oxide or agriculturally-suitable salt thereof.

19. A fungicidal composition comprising an effective amount of a compound of Formula I, II, or III:



wherein:

n is 0, 1 or 2;

Q is independently O or S;

R^1 is C_3 - C_{10} alkyl; C_3 - C_5 cycloalkyl; C_4 - C_{10} alkenyl; C_4 - C_{10} alkynyl; C_1 - C_{10} haloalkyl; C_3 - C_{10} haloalkenyl; C_3 - C_{10} haloalkynyl; C_2 - C_{10} alkoxyalkyl; C_2 - C_{10} alkylthioalkyl; C_2 - C_{10} alkylsulfinylalkyl; C_2 - C_{10} alkylsulfonylalkyl; C_5 - C_{10} cycloalkylalkyl; C_4 - C_{10} alkenyloxyalkyl; C_4 - C_{10} alkynyloxyalkyl; C_4 - C_{10} (cycloalkyl)oxyalkyl; C_4 - C_{10} alkenylthioalkyl; C_4 - C_{10} alkynylthioalkyl; C_6 - C_{10} (cycloalkyl)thioalkyl; C_2 - C_{10} haloalkoxyalkyl; C_4 - C_{10} haloalkenyloxyalkyl; C_4 - C_{10} haloalkynyloxyalkyl; C_4 - C_{10} alkoxyalkenyl; C_4 - C_{10} alkoxyalkynyl; C_4 - C_{10} alkylthioalkenyl; C_4 - C_{10} alkylthioalkynyl; C_4 - C_{10} trialkylsilylalkyl; C_1 - C_{10} alkyl substituted with $NR^{11}R^{12}$, nitro, cyano, or phenyl optionally substituted with R^{14} , R^{15} , and R^{16} ; C_1 - C_{10} alkoxy; C_1 - C_{10} haloalkoxy; C_1 - C_{10} alkylthio; C_1 - C_{10} haloalkylthio; $NR^{11}R^{12}$; or pyridyl, furanyl, thienyl, naphthyl, benzofuranyl,

benzothienyl, or quinolinyll each optionally substituted with R¹⁴, R¹⁵, and R¹⁶;

R² is C₃-C₁₀ alkyl; C₆-C₇ cycloalkyl; C₃-C₁₀ alkenyl; C₃-C₁₀ alkynyl; C₁-C₁₀ haloalkyl; C₃-C₁₀ haloalkenyl; C₃-C₁₀ haloalkynyl; C₂-C₁₀ alkoxyalkyl; C₂-C₁₀ alkylthioalkyl; C₂-C₁₀ alkylsulfinylalkyl; C₂-C₁₀ alkylsulfonylalkyl; C₄-C₁₀ cycloalkylalkyl; C₄-C₁₀ alkenyloxyalkyl; C₄-C₁₀ alkynyloxyalkyl; C₄-C₁₀ (cycloalkyl)oxyalkyl; C₄-C₁₀ alkenylthioalkyl; C₄-C₁₀ alkynylthioalkyl; C₆-C₁₀ (cycloalkyl)thioalkyl; C₂-C₁₀ haloalkoxyalkyl; C₄-C₁₀ haloalkenyloxyalkyl; C₄-C₁₀ haloalkynyloxyalkyl; C₄-C₁₀ alkoxyalkenyl; C₄-C₁₀ alkoxyalkynyl; C₄-C₁₀ alkylthioalkenyl; C₄-C₁₀ alkylthioalkynyl; C₄-C₁₀ trialkylsilylalkyl; C₃-C₁₀ cyanoalkyl; C₂-C₁₀ nitroalkyl; C₁-C₁₀ alkyl substituted with CO₂R¹¹, NR¹¹R¹², or phenyl optionally substituted with R¹³, R¹⁵, and R¹⁶; phenyl optionally substituted with R¹³, R¹⁵, and R¹⁶; -N=CR¹¹R¹¹; or -NR¹¹R¹²; or

R¹ and R² are taken together to form -CH₂(CH₂)_mCH₂-; m is 1-4;

R³ is halogen; C₁-C₈ alkyl; C₃-C₈ cycloalkyl; C₂-C₈ alkenyl; C₂-C₈ alkynyl; C₁-C₈ haloalkyl; C₃-C₈ haloalkenyl; C₃-C₈ haloalkynyl; C₁-C₈ alkoxy; C₁-C₈ haloalkoxy; C₃-C₈ alkenyloxy; C₃-C₈ alkynyloxy; C₁-C₈ alkylthio; C₃-C₈ alkenylthio; C₃-C₈ alkynylthio; C₁-C₈ alkylsulfinyl; C₁-C₈ alkylsulfonyl; C₂-C₈ alkoxyalkyl; C₂-C₈ alkylthioalkyl; C₂-C₈ alkylsulfinylalkyl; C₂-C₈ alkylsulfonylalkyl; C₄-C₈ cycloalkylalkyl; C₃-C₈ trialkylsilyl; nitro; NR¹¹R¹²; C₅-C₈ trialkylsilylalkynyl; or phenyl optionally substituted with at least one R¹³;

R⁴ is hydrogen; halogen; C₁-C₄ alkyl; C₁-C₄ haloalkyl; C₁-C₄ alkoxy; or C₁-C₄ haloalkoxy;

R⁵ is C₃-C₅ alkyl; C₇-C₁₀ alkyl; C₄-C₇ alkenyl; C₃-C₅ alkynyl; C₁-C₁₀ haloalkyl; C₅-C₁₀ haloalkenyl; C₃-C₁₀ haloalkynyl; C₂-C₁₀ alkoxyalkyl other than methoxypropyl; C₂-C₁₀ alkylthioalkyl; C₂-C₁₀ alkylsulfinylalkyl; C₂-C₁₀ alkylsulfonylalkyl; C₄-C₁₀ cycloalkylalkyl; C₄-C₁₀ alkenyloxyalkyl; C₄-C₁₀ alkynyloxyalkyl; C₄-C₁₀ (cycloalkyl)oxyalkyl; C₄-C₁₀ alkenylthioalkyl; C₄-C₁₀ alkynylthioalkyl; C₆-C₁₀ (cycloalkyl)thioalkyl; C₂-C₁₀ haloalkoxyalkyl; C₄-C₁₀ haloalkenyloxyalkyl; C₄-C₁₀ haloalkynyloxyalkyl; C₄-C₁₀ alkoxyalkenyl; C₄-C₁₀ alkoxyalkynyl; C₄-C₁₀ alkylthioalkenyl; C₄-C₁₀ alkylthioalkynyl; C₄-C₁₀ trialkylsilylalkyl; C₁-C₁₀ alkyl substituted with NR¹¹R¹², nitro, or phenyl optionally substituted with at least one of R¹⁴, R¹⁵, and R¹⁶; C₂-C₁₀ alkyl substituted with cyano; C₁-C₁₀ alkoxy; C₁-C₁₀ haloalkoxy; C₁-C₁₀ alkylthio; C₁-C₁₀ haloalkylthio; NR¹¹R¹²; or

phenyl, furanyl, thienyl, naphthyl, benzofuranyl, or benzothienyl each optionally substituted with R¹⁴, R¹⁵, and R¹⁶;

R⁶ is C₃-C₁₀ alkyl; C₃-C₇ alkenyl; C₃-C₁₀ alkynyl; C₁-C₁₀ haloalkyl; C₃-C₁₀ haloalkenyl; C₃-C₁₀ haloalkynyl; C₃-C₁₀ alkoxyalkyl other than propoxymethyl; C₂-C₁₀ alkylthioalkyl; C₂-C₁₀ alkylsulfinylalkyl; C₂-C₁₀ alkylsulfonylalkyl; C₄-C₁₀ cycloalkylalkyl; C₄-C₁₀ alkenyloxyalkyl; C₄-C₁₀ alkynyloxyalkyl; C₄-C₁₀ (cycloalkyl)oxyalkyl; C₄-C₁₀ alkenylthioalkyl; C₄-C₁₀ alkynylthioalkyl; C₆-C₁₀ (cycloalkyl)thioalkyl; C₂-C₁₀ haloalkoxyalkyl; C₄-C₁₀ haloalkenyloxyalkyl; C₄-C₁₀ haloalkynyloxyalkyl; C₄-C₁₀ alkoxyalkenyl; C₄-C₁₀ alkoxyalkynyl; C₄-C₁₀ alkylthioalkenyl; C₄-C₁₀ alkylthioalkynyl; C₄-C₁₀ trialkylsilylalkyl; C₅-C₁₀ cyanoalkyl; C₂-C₁₀ nitroalkyl; or C₃-C₁₀ alkyl substituted with CO₂R¹¹, NR¹¹R¹², or phenyl optionally substituted with R¹³, R¹⁵, and R¹⁶; or phenyl optionally substituted with R¹³, R¹⁵, and R¹⁶; or

R⁵ and R⁶ are taken together to form -CH₂(CH₂)_mCH₂-;

R⁷ is C₃-C₁₀ alkyl; C₃-C₇ cycloalkyl; C₄-C₇ alkenyl; propynyl; C₅-C₁₀ alkynyl; C₂-C₁₀ haloalkyl; C₃-C₁₀ haloalkenyl; C₃-C₁₀ haloalkynyl; C₂-C₁₀ alkoxyalkyl; C₂-C₁₀ alkylthioalkyl; C₂-C₁₀ alkylsulfinylalkyl; C₂-C₁₀ alkylsulfonylalkyl; C₄-C₁₀ alkenyloxyalkyl; C₄-C₁₀ alkynyloxyalkyl; C₄-C₁₀ (cycloalkyl)oxyalkyl; C₄-C₁₀ alkenylthioalkyl; C₄-C₁₀ alkynylthioalkyl; C₆-C₁₀ (cycloalkyl)thioalkyl; C₂-C₁₀ haloalkoxyalkyl; C₄-C₁₀ haloalkenyloxyalkyl; C₄-C₁₀ haloalkynyloxyalkyl; C₄-C₁₀ alkoxyalkenyl; C₄-C₁₀ alkoxyalkynyl; C₄-C₁₀ alkylthioalkenyl; C₄-C₁₀ alkylthioalkynyl; C₄-C₁₀ trialkylsilylalkyl; C₁-C₁₀ alkyl substituted with NR¹¹R¹² or nitro; C₂-C₁₀ alkyl substituted with cyano; C₁-C₁₀ alkoxy; C₁-C₁₀ haloalkoxy; C₁-C₁₀ alkylthio; C₁-C₁₀ haloalkylthio; NR¹²R¹⁷; or phenyl, pyridyl, furanyl, thienyl, naphthyl, benzofuranyl, benzothienyl, or quinolinyl each optionally substituted with R¹⁴, R¹⁵, and R¹⁶;

R⁸ is hydrogen; C₁-C₄ alkyl; or -C(=O)R¹⁰;

R⁹ is hydrogen; C₂-C₁₀ alkyl; C₃-C₇ cycloalkyl; C₃-C₁₀ alkenyl; C₃-C₁₀ alkynyl; C₃-C₁₀ haloalkyl; C₃-C₁₀ haloalkenyl; C₃-C₁₀ haloalkynyl; C₃-C₁₀ alkoxyalkyl other than butoxyethyl; C₂-C₁₀ alkylthioalkyl; C₂-C₁₀ alkylsulfinylalkyl; C₂-C₁₀ alkylsulfonylalkyl; C₄-C₁₀ cycloalkylalkyl; C₄-C₁₀ alkenyloxyalkyl; C₄-C₁₀ alkynyloxyalkyl; C₄-C₁₀ (cycloalkyl)oxyalkyl; C₄-C₁₀ alkenylthioalkyl; C₄-C₁₀ alkynylthioalkyl; C₆-C₁₀ (cycloalkyl)thioalkyl; C₂-C₁₀ haloalkoxyalkyl; C₄-C₁₀ haloalkenyloxyalkyl; C₄-C₁₀ haloalkynyloxyalkyl; C₄-C₁₀ alkoxyalkenyl; C₄-C₁₀ alkoxyalkynyl; C₄-C₁₀ alkylthioalkenyl; C₄-C₁₀ alkylthioalkynyl; C₄-C₁₀ trialkylsilylalkyl;

- C₁-C₁₀ alkyl substituted with NR¹¹R¹²; C₄-C₁₀ cyanoalkyl; C₂-C₁₀ nitroalkyl; C₁-C₈ alkyl substituted with CO₂R¹¹; pyridyl, furanyl, thienyl, or naphthyl each optionally substituted with R¹⁴, R¹⁵, and R¹⁶; -N=CR¹¹R¹¹; -NR¹²R¹⁷; -OR¹²; or -NC(=Q)NR¹¹R¹²; or R³ and R⁴ are both iodine and R⁹ is phenyl optionally substituted with R¹⁴, R¹⁵, and R¹⁶; or
 5 R⁷ and R⁹ are taken together to form -CH₂(CH₂)_mCH₂-;
 R¹⁰ is hydrogen; C₁-C₄ alkyl; C₁-C₄ alkoxy; or NR¹¹R¹²;
 R¹¹ is independently hydrogen; C₁-C₄ alkyl; or phenyl optionally substituted with at least one R¹³;
 10 R¹² is independently hydrogen; C₁-C₈ alkyl; or phenyl optionally substituted with at least one R¹³; or
 R¹¹ and R¹² are taken together to form -CH₂CH₂CH₂CH₂-, -CH₂(CH₂)₃CH₂-, -CH₂CH₂OCH₂CH₂-, -CH₂CH(Me)CH₂CH(Me)CH₂-, or -CH₂CH(Me)OCH(Me)CH₂-;
 15 R¹³ is independently halogen; C₁-C₄ alkyl; C₁-C₄ alkoxy; C₁-C₄ haloalkyl; nitro; or cyano;
 R¹⁴ is independently C₁-C₆ alkyl; C₁-C₆ alkoxy; C₁-C₆ haloalkyl; halogen; C₂-C₈ alkynyl; C₁-C₆ thioalkyl; phenyl or phenoxy each optionally substituted with at least one R¹³; cyano; nitro; C₁-C₆ haloalkoxy; C₁-C₆ haloalkylthio; C₂-C₆
 20 alkenyl; C₂-C₆ haloalkenyl; acetyl; CO₂Me; or N(C₁-C₂ alkyl)₂;
 R¹⁵ is independently methyl; ethyl; methoxy; methylthio; halogen; or trifluoromethyl;
 R¹⁶ is independently halogen; and
 R¹⁷ is independently C₁-C₈ alkyl; or phenyl optionally substituted with at least one
 25 R¹³; or
 an *N*-oxide or agriculturally-suitable salt thereof; and
 at least one of (a) a surfactant, (b) an organic solvent, and (c) at least one solid or liquid diluent.

- 30 20. A method for controlling wheat powdery mildew comprising applying to the plant or portion thereof to be protected, to the media in which the plant to be protected is growing, or to the plant seed or seedling to be protected an effective amount of a compound of Claim 2.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 94/04965

A. CLASSIFICATION OF SUBJECT MATTER

IPC 5 C07D239/95 C07D239/96 C07D401/04 C07D405/04 C07D409/04
A01N43/54

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 5 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 276 825 (NISSHIN FLOUR MILLING) 3 August 1988 see claims ---	1,3
X	US,A,3 755 582 (G.A.BULLOCK ET AL.) 28 August 1973 cited in the application see the whole document ---	1-3, 13-15
X	US,A,3 784 693 (D.HABECK ET AL.) 8 January 1974 see column 1 - column 8 ---	1,2
X	US,A,3 867 384 (G. A. BULLOCK) 18 February 1975 cited in the application see the whole document ---	1-3, 13-15
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "I" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

15 September 1994

Date of mailing of the international search report

26.09.94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Francois, J

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 115, no. 7, 1991, Columbus, Ohio, US; abstract no. 71634p, page 803 ; see abstract & DD,A,287 033 (AKAD. DER WISSENSCHAFTEN) 14 February 1991 ---	1,2,13
X	CHEMICAL ABSTRACTS, vol. 89, no. 19, 1978, Columbus, Ohio, US; abstract no. 163530z, P.BHARGAVA ET AL. 'SYNTHESIS AND SPECTRAL STUDIES ON SOME TYPICAL 4(3H)QUINAZOLONES AS POSSIBLE ANTIMALARIALS.' page 573 ; see abstract & EGYPT. J. CHEM., vol.18, no.3, 1975, CAIRO pages 393 - 401 ---	1,3,13
X	CHEMICAL ABSTRACTS, vol. 89, no. 13, 1978, Columbus, Ohio, US; abstract no. 109345e, P.BHARGAVA ET AL. 'SYNTHESIS OF SOME NEW 6 ,8-DIIODO-S-SUBSTITUTED-2-THIO-3-ARYL-4(3H))QUINAZOLINONE' page 908 ; see abstract & J.SCI.RES.BANARAS HINDU UNIV., vol.26, no.1, 1976, INDIA pages 27 - 32 ---	1,3
X	CHEMICAL ABSTRACTS, vol. 85, no. 1, 1976, Columbus, Ohio, US; abstract no. 5582f, P.BHARGAVA ET AL. 'STUDIES ON 4(3H)QUINAZOLONES.' page 450 ; see abstract & J.INDIAN CHEM. SOC., vol.53, no.1, 1976, INDIA pages 46 - 49 ---	1,3
X	CHEMICAL ABSTRACTS, vol. 107, no. 5, 1987, Columbus, Ohio, US; abstract no. 36462u, R.LAKHAN ET AL. 'SYNTHESIS AND ANTIBACTERIAL ACTIVITY OF SOME 4-(3H)-QUINAZOLINONES.' page 396 ; see abstract & J.CHEM.ENG.DATA, vol.32, no.3, 1987, INDIA pages 384 - 386 -----	1,3,13

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US 94/04965

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0276825	03-08-88	JP-A- 63295565	01-12-88
		DE-A- 3874358	15-10-92
		ES-T- 2044981	16-01-94
		US-A- 4861780	29-08-89
		US-A- 5008266	16-04-91
US-A-3755582	28-08-73	NONE	
US-A-3784693	08-01-74	NONE	
US-A-3867384	18-02-75	NONE	
DD-A-287033		NONE	

SEARCHED BY: [illegible] INDEXED BY: [illegible]

THIS PAGE BLANK (USPTO)